

A Characterization of Ternary Rings of Operators*

HEINRICH ZETTL[†]*Fachbereich Mathematik, Universität des Saarlandes,
D-6600 Saarbrücken, Federal Republic of Germany*

0. INTRODUCTION

A *ternary ring of operators* (TRO) between two complex Hilbert spaces H and K is a linear subspace \mathfrak{R} of $\mathcal{L}(H, K)$ satisfying $AB^*C \in \mathfrak{R}$ for all $A, B, C \in \mathfrak{R}$. The purpose of this paper is to give a characterization of a TRO as an abstract ternary ring.

In [2] Hestenes introduced a purely algebraic concept of an abstract ternary ring which was related to the study of TROs between finite dimensional spaces and which was pursued further by other authors, e.g., [10, 11]. Our approach applies to infinite dimensional spaces and is oriented at the representation of a C^* -algebra (respectively, W^* -algebra) as a norm (respectively, weakly) closed star algebra of operators on a Hilbert space. Thus, we confine our considerations to TROs which are norm or weakly closed. The first step towards a characterization of these rings is to find a suitable concept of an abstract ternary ring which reflects the ternary operation, $(A, B, C) \mapsto AB^*C$, the norm relations, and, in the second case, also the weak-operator structure. This leads us to the following basic definition:

0.1. DEFINITION. A *ternary C^* -ring* $(\mathfrak{X}, (\cdot, \cdot, \cdot), \|\cdot\|)$ consists of a complex Banach space $(\mathfrak{X}, \|\cdot\|)$ and a ternary operation $(\cdot, \cdot, \cdot): \mathfrak{X} \times \mathfrak{X} \times \mathfrak{X} \rightarrow \mathfrak{X}$ such that for all $v, w, x, y, z \in \mathfrak{X}$ and $\lambda \in \mathbb{C}$ the following conditions hold:

$$(\lambda x + y, v, w) = \lambda(x, v, w) + (y, v, w), \quad (0.1)$$

$$(v, w, \lambda x + y) = \lambda(v, w, x) + (v, w, y),$$

$$(v, \lambda x + y, w) = \bar{\lambda}(v, x, w) + (v, y, w), \quad (0.2)$$

$$((v, w, x), y, z) = (v, (y, x, w), z) = (v, w, (x, y, z)), \quad (0.3)$$

* This article is based on the author's doctoral thesis written at the university of Erlangen under the supervision of Professor D. Kölzow.

[†] Current address: Mathematisches Institut der Universität Erlangen, Bismarckstr. 1 1/2, D-8520 Erlangen, Federal Republic of Germany.

$$\|(x, y, z)\| \leq \|x\| \|y\| \|z\|, \quad (0.4)$$

$$\|(x, x, x)\| = \|x\|^3. \quad (0.5)$$

A *ternary W^* -ring* is a ternary C^* -ring which is, in addition, a dual space.

Starting from these definitions we prove the following representation theorems which we give here in shortened versions.

THEOREM 3. *For each ternary C^* -ring $(\mathfrak{X}, (\cdot, \cdot, \cdot), \|\cdot\|)$ there exists one, and only one, operator $T: \mathfrak{X} \rightarrow \mathfrak{X}$ satisfying*

$$(1) \quad T^2 = I,$$

$$(2) \quad T((x, y, z)) = (Tx, y, z) = (x, Ty, z) = (x, y, Tz) \text{ for all } x, y, z \in \mathfrak{X},$$

and

$$(3) \quad (\mathfrak{X}, T \circ (\cdot, \cdot, \cdot), \|\cdot\|) \text{ is a ternary } C^*\text{-ring which is isomorphic to a norm closed TRO.}$$

THEOREM 4.1. *For each ternary W^* -ring $(\mathfrak{X}, (\cdot, \cdot, \cdot), \|\cdot\|)$ the operator T is weak* continuous, and $(\mathfrak{X}, T \circ (\cdot, \cdot, \cdot), \|\cdot\|)$ is a ternary W^* -ring which is (isometrically and weakly*) isomorphic to a weakly closed TRO.*

These results are based upon two representation theorems for Hilbert modules over C^* -algebras which say, first, that each such module can be faithfully represented as a norm closed TRO (cf. Theorem 2.6), and, second, that a self-dual Hilbert module over a W^* -algebra is isomorphic to a weakly closed TRO (cf. Theorem 2.8). For the proof in both cases we use a GNS-construction extending ideas found in [5].

The deduction of our main theorems from these results is as follows: We construct a C^* -algebra $\mathfrak{A} \subset \mathcal{L}(\mathfrak{X})$ and a form $\alpha: \mathfrak{X} \times \mathfrak{X} \rightarrow \mathfrak{A}$ which contain all information on $(\mathfrak{X}, (\cdot, \cdot, \cdot), \|\cdot\|)$, in the sense that

$$(x, y, z) = \alpha(z, y)(x) \quad \text{and} \quad \|x\|^2 = \|\alpha(x, x)\|$$

hold for all $x, y, z \in \mathfrak{X}$ (cf. Proposition 3.2). Moreover, the pair (\mathfrak{X}, α) satisfies the axioms for a Hilbert \mathfrak{A} -module, apart from α being positive. If we set $T = 2P - I$, where $P \in \mathcal{L}(\mathfrak{X})$ is the projection with range $\{x \mid \alpha(x, x) \geq 0\}$ and kernel $\{x \mid \alpha(x, x) \leq 0\}$, we see that the form $\alpha_T: (x, y) \mapsto \alpha(Tx, y)$ is positive and that (\mathfrak{X}, α_T) is a Hilbert \mathfrak{A} -module on which then 2.6 can be applied (cf. 3.7 infra). In the case of a ternary W^* -ring the double centralizer algebra $M(\mathfrak{A})$ of \mathfrak{A} turns out to be a W^* -algebra, and (\mathfrak{X}, α_T) becomes a self-dual Hilbert $M(\mathfrak{A})$ -module (cf. Corollary 4.10), so that Theorem 4.1 is a consequence of Theorem 2.8.

We then prove that $M(\mathfrak{A})_*$ is a quotient of $\mathfrak{X} \hat{\otimes}_\gamma \mathfrak{X}_*$ (cf. Corollary 4.11), and show that the predual \mathfrak{X}_* of \mathfrak{X} coincides with $\mathfrak{X}_c \otimes M(\mathfrak{A})_*$ (cf. Example

4.12). Since \mathfrak{A} depends only on \mathfrak{X} , the predual is uniquely determined up to isomorphism. Finally, we prove that each weakly closed TRO is a self-dual Hilbert module over a W^* -algebra, and thus a ternary W^* -ring (cf. 4.12).

On the other hand, it is fairly obvious that any norm closed TRO between two Hilbert spaces (such as the space of all compact operators or a C^* -algebra in $\mathcal{L}(H)$) is a ternary C^* -ring. All these examples lead to $T = I$; in general, however, $T \neq I$ (cf. Example 3.8). Hence, the problem is to characterize the case $T = I$. We have no answer to this question, but give two ideas to it at the end of Section 3.

In our considerations we need some fundamental properties of TROs: in particular we need statements concerning partial isometries and a density theorem of Kaplansky type. All this is provided in Section 1.

We make the following conventions: The linear spaces considered here are over the complex field, \mathbb{C} . If X is a linear space, then X_c denotes X equipped with the conjugate scalar multiplication $(\lambda, x) \mapsto \bar{\lambda}x$, and we write I for the identity operator on X . If X and Y are normed spaces, then $\mathcal{L}(X, Y)$ denotes the set of all bounded operators from X to Y . The symbol X^* (respectively, X_*) is used for the dual space of X (respectively, predual of X whenever it exists), and $\bar{\Omega}^n$ stands for the norm closure of $\Omega \subset X$. The action of an algebra \mathfrak{A} on a right \mathfrak{A} -module X is denoted by $(x, a) \mapsto x \cdot a$, and it is always assumed to satisfy $\lambda(x \cdot a) = (\lambda x) \cdot a = x \cdot (\lambda a)$ for all $\lambda \in \mathbb{C}$, $x \in X$, $a \in \mathfrak{A}$. Similarly, for left \mathfrak{A} -modules. The reference for Banach \mathfrak{A} -modules and approximate identities is [13, Part I, Sect. 2]. The notions and facts on C^* -algebras and W^* -algebras we use can be found in [9].

1. TERNARY RINGS OF OPERATORS: PRELIMINARIES

This section collects some basic material which will subsequently be used in connection with TROs arising from Representation Theorems 2.6 and 3.1.

In the following, \mathfrak{K} denotes a norm closed TRO between two Hilbert spaces H and K . It is clear that the weak-operator closure $\bar{\mathfrak{K}}^\sigma$ of \mathfrak{K} is also a TRO. By $[\mathfrak{K}]_H$ (respectively, $\{\mathfrak{K}\}_H$) we shall denote the C^* -algebra (respectively, W^* -algebra) generated by $\{A^*B \mid A, B \in \mathfrak{K}\}$ in $\mathcal{L}(H)$. Similarly, we define $[\mathfrak{K}]_K$ and $\{\mathfrak{K}\}_K$.

For instance, the space $\mathfrak{C}(H, K)$ of all compact operators is a norm closed TRO with weak closure $\mathcal{L}(H, K)$, while $[\mathfrak{C}(H, K)]_H$ is equal to $\mathfrak{C}(H)$. When \mathfrak{A} is a C^* -algebra and $\{\pi, H\}$, $\{\pi', K\}$ are two $*$ -representations of \mathfrak{A} (cf. [9, Def. 1.16.3]), the intertwining ring $\mathfrak{I} = \{A \in \mathcal{L}(H, K) \mid A\pi(a) = \pi'(a)A, a \in \mathfrak{A}\}$ is a weakly closed TRO, and $\{\mathfrak{I}\}_H$ coincides with the commutant of $\pi(\mathfrak{A})$ in $\mathcal{L}(H)$.

Our first considerations are related to the partial isometries in $\bar{\mathfrak{K}}^\sigma$. These appear naturally in the polar decomposition of operators (cf.

Proposition 1.2) and provide the extreme points in the unit ball of $\overline{\mathfrak{R}}^\sigma$ (cf. Lemma 1.3). The latter fact will lead to a polar decomposition for weak* continuous functionals on a ternary W^* -ring (cf. Lemma 4.4). Further, we shall make essential use of C^* -algebras associated with a single partial isometry. To this end we introduce the following notations which are partly due to [3, Sect. 15]:

1.1. *Notations.* Let $R \in \mathfrak{R}$ be a partial isometry, i.e., $R = RR^*R$, and denote by E and F the support and range projection R^*R and RR^* of R , respectively. Define

$$\mathfrak{R}_{(R)} = \{X \in \mathfrak{R} \mid X = XE = FX\}.$$

Then, endowed with the operator norm, $\mathfrak{R}_{(R)}$ becomes a C^* -algebra with unit R by

$$X \circ Y = XR^*Y, \quad X^\# = RX^*R. \quad (1.1)$$

Obviously, $X \mapsto R^*X$ is a $*$ -isomorphism of $\mathfrak{R}_{(R)}$ onto $E|\mathfrak{R}|_H E$.

Next, put $\mathfrak{S} = \overline{\mathfrak{R}}^\sigma$, so that $\mathfrak{S}_{(R)} = \overline{\mathfrak{R}_{(R)}}^\sigma$. Then $\mathfrak{S}_{(R)}$ is a W^* -algebra since the $*$ -isomorphism $X \mapsto R^*X$ of $\mathfrak{S}_{(R)}$ onto $E|\mathfrak{R}|_H E$ is a homeomorphism in the weak-operator topologies. Note that $\mathfrak{R}_{(R)}$ (respectively, $\mathfrak{S}_{(R)}$) is a C^* -subalgebra (respectively, W^* -subalgebra) of $\mathcal{L}(H, K)_{(R)}$.

1.2. PROPOSITION. Let $X \in \mathfrak{R}$. If R denotes the partial isometry of the polar decomposition $X = R|X|$ of X , then the C^* -algebra (respectively, W^* -algebra) generated by X in $\mathcal{L}(H, K)_{(R)}$ is equal to the smallest norm (respectively, weakly) closed TRO containing X . In particular, $R \in \overline{\mathfrak{R}}^\sigma$.

Proof. Let $P = \mathbb{C}[t]$ denote the polynomials, define $P_0 = \{p \in P \mid p(0) = 0\}$ and write $p_R(X)$ for the polynomial associated with X and $p \in P$ in $\mathcal{L}(H, K)_{(R)}$. By [3, Theorem 6.2], $|X| = R^*X$. Since $A \mapsto RA$ is the inverse of the $*$ -isomorphism $Y \mapsto R^*Y$ of $\mathcal{L}(H, K)_{(R)}$ onto $R^*R\mathcal{L}(H)R^*R$ we obtain, using the first equation in (1.1) for $\mathcal{L}(H, K)$ instead of \mathfrak{R} ,

$$Xp(|X|) = X \circ (Rp(|X|)) = X \circ p_R(X) = (tp)_R(X).$$

Hence, the C^* -algebra $\mathfrak{C} = \overline{\{p_R(X) \mid p \in P_0\}}^n$ generated by X in $\mathcal{L}(H, K)_{(R)}$ is equal to $\overline{\{Xp(|X|) \mid p \in P\}}^n$.

Now if \mathfrak{T} denotes the norm closed TRO generated by X , then $|X| \in [\mathfrak{T}]_H$ and $\mathfrak{T}([\mathfrak{T}]_H \oplus \mathbb{C}) \subset \mathfrak{T}$, which imply $\mathfrak{C} \subset \mathfrak{T}$. On the other hand, by (1.1),

$$p_R(X)q_R(X)^*r_R(X) = p_R(X) \circ q_R(X)^\# \circ r_R(X) = (p\bar{q}r)_R(X).$$

So, \mathfrak{C} is a norm closed TRO containing X . Thus, $\mathfrak{C} = \mathfrak{T}$.

Similarly, $\overline{\{p_R(X) \mid p \in P_0\}}^\sigma = \overline{\mathfrak{T}}^\sigma$. The unit S of the W^* -subalgebra $\overline{\mathfrak{T}}^\sigma$ of

$\mathcal{L}(H, K)_{(R)}$ is a partial isometry in $\mathcal{L}(H, K)$, because $S = S \circ S^\# \circ S = SR^*RS^*RR^*S = SS^*S$. Moreover,

$$SS^*X = SR^*RS^*RR^*X = S \circ S^\# \circ X = S \circ X = X$$

and, analogously, $XS^*S = X$. Thus, $SS^* \geq RR^*$ and $S^*S \geq R^*R$ in $\mathcal{L}(K)$ and $\mathcal{L}(H)$, respectively, and therefore $S = R$. In particular, we have $R \in \mathfrak{T}^\sigma \subset \mathfrak{R}^\sigma$.

1.3. LEMMA. *For each X in the unit ball $D_1(\mathfrak{R})$ of \mathfrak{R} the following conditions are equivalent:*

- (1) X is an extreme point of $D_1(\mathfrak{R})$;
- (2) X is a partial isometry satisfying $(I - XX^*)\mathfrak{R}(I - X^*X) = \{0\}$;
- (3) $(I - XX^*)\mathfrak{R}(I - X^*X) = \{0\}$.

Proof. Let E and F denote the support and range projection of X , respectively. Observe that

$$(I - F)\mathfrak{R}(I - E) = \{Y \in \mathfrak{R} \mid 0 = Y^*X, 0 = YX^*\}. \quad (1.2)$$

(1) \Rightarrow (2). Let R be the partial isometry of the polar decomposition $X = R|X|$ of X . In view of 1.1 and 1.2, $Y \mapsto R^*Y$ is a $*$ -isomorphism of the norm closed TRO generated by X onto the C^* -algebra generated by $|X| = R^*X$. Thus, using [9, Proposition 1.6.2], X is a partial isometry. Moreover, by (1.2), for each $Y \in (I - F)\mathfrak{R}(I - E)$ with $\|Y\| \leq 1$ we have

$$\|Y + X\|^2 = \|(Y + X)^*(Y + X)\| = \|Y^*Y + X^*X\| = \text{Max}\{\|X\|^2, \|Y\|^2\} \leq 1$$

which yields $Y = 0$, as desired.

(3) \Rightarrow (1). From $0 = (I - XX^*)X(I - X^*X)$ we obtain $0 = X^*X(I - X^*X)$. Hence, X^*X is a projection, and so, X is a partial isometry. Now, fix $Y, Z \in D_1(\mathfrak{R})$ with $X = (Y + Z)/2$. Since FYE and FZE belong to the unit ball of the C^* -algebra $\mathfrak{R}_{(X)}$, the arguments in [9, Theorem 1.6.4] show that $X = FYE = FZE$. From

$$\begin{aligned} 1 &\geq \|YE\|^2 = (FYE + (I - F)YE)^*(FYE + (I - F)YE)\| \\ &= \|X + (I - F)YE\|^2 = 1 + \|(I - F)YE\|^2 \end{aligned}$$

we deduce that $FYE = YE$. Similarly, $FZE = ZE$. Thus, $X = YE = ZE$.

Since (3) implies that $(I - E)\mathfrak{R}^*(I - F) = \{0\}$, the same line of reasoning with X^* and \mathfrak{R}^* instead of X and \mathfrak{R} yields $X = FY = FZ$. Therefore,

$$\begin{aligned} Y &= YE + Y(I - E) \\ &= X + FY(I - E) + (I - F)Y(I - E) = X \quad \text{and} \quad Z = X. \end{aligned}$$

Our next result is an analogue of Kaplansky's density theorem which we shall use for Hilbert modules and ternary W^* -rings. For the proof we modify the arguments in [13, Sect. 4, Theorem 2.5].

1.4. PROPOSITION. *The unit ball of \mathfrak{K} is weakly and strongly dense in the unit ball of the weak closure $\tilde{\mathfrak{K}}^\sigma$ of \mathfrak{K} .*

Proof. Obviously,

$$\tilde{\mathfrak{K}} = \left\{ \begin{pmatrix} 0 & Y^* \\ X & 0 \end{pmatrix} \mid X, Y \in \mathfrak{K} \right\}$$

is a norm closed, self-adjoint TRO in $\mathcal{L}(H \oplus K)$, and

$$X \mapsto \begin{pmatrix} 0 & X^* \\ X & 0 \end{pmatrix}$$

is a real-linear isometry and a weak homeomorphism of \mathfrak{K} onto the self-adjoint part $\tilde{\mathfrak{K}}_{sa}$ of $\tilde{\mathfrak{K}}$. Moreover, observe that the weak and strong operator topologies coincide on convex sets. Thus, if we assume that $H = K$ and that \mathfrak{K} is self-adjoint, it remains to show that $D_1(\mathfrak{B})$ is contained in the strong closure of $D_1(\mathfrak{A})$, where $\mathfrak{A} = \mathfrak{K}_{sa}$, $\mathfrak{B} = (\mathfrak{K}^\sigma)_{sa}$. To this end we need

LEMMA 1.4.1. *Let $X \in \mathcal{L}(H)_{sa}$ and let \mathfrak{T} denote the norm closed TRO generated by X . Then there is an $X' \in \mathfrak{T}$ satisfying $X = 2X'(I + X'^2)^{-1}$.*

Proof. By 1.2, \mathfrak{T} is equal to the C^* -algebra generated by X in $\mathcal{L}(H)_{(R)}$. Let $\chi \mapsto \chi_R(X)$ denote its functional calculus. Define the function f on $[0, 1]$ by $f(t) = 2t/(1 + t^2)$, and write $X' = (f^{-1})_R(X)$. Then X' is positive in the C^* -algebra \mathfrak{T} , and thus R^*X' is positive in $\mathcal{L}(H)$. Since $X = X^*$ there are two projections E and F with $R = E - F$, $EF = 0$. From $RX = XR$ we conclude that \mathfrak{T} commutes with R in $\mathcal{L}(H)$. Hence, $RX' = X'R$. Moreover, $X' = X'^*$, as $X'R$ is positive. Therefore,

$$X = f_R(X') = Rf(RX') = R(2RX'(I + R^2X'^2)^{-1}) = 2X'(I + X'^2)^{-1},$$

which proves Lemma 1.4.1.

Now, fix $X \in D_1(\mathfrak{B})$, write $X = 2X'(I + X'^2)^{-1}$ with $X' \in \mathfrak{B}$ according to Lemma 1.4.1, and define $Y = 2Y'(I + Y'^2)^{-1}$, $Y' \in \mathfrak{A}$. Then, for each $Y' \in \mathfrak{A}$ we have $\|Y\| \leq 1$, $\|(I + Y'^2)^{-1}\| \leq 1$ and

$$(X - Y)/2 = (I + Y'^2)^{-1}((X' - Y') + Y'(Y' - X')X')(I + X'^2)^{-1}.$$

Since \mathfrak{B} is contained in the strong closure of \mathfrak{A} we are done.

The following statement is necessary for the proof of Corollary 4.7:

1.5. LEMMA. *If \mathfrak{I} is a weakly closed two-sided ideal in $\bar{\mathfrak{R}}^\sigma$ (that is, a subspace of $\bar{\mathfrak{R}}^\sigma$ satisfying $\bar{\mathfrak{I}}^\sigma = \mathfrak{I} \supset \mathfrak{I}\mathfrak{R}^*\mathfrak{R} \cup \mathfrak{R}\mathfrak{R}^*\mathfrak{I}$), then there is a central projection $E \in \{\mathfrak{R}\}_H$ such that $\mathfrak{I} = \bar{\mathfrak{R}}^\sigma E$.*

Proof. Put $\mathfrak{U} = \{\mathfrak{R}\}_H$ and $\mathfrak{B} = \{\mathfrak{I}\}_H$. Then $\mathfrak{I}\mathfrak{U} \subset \mathfrak{I}$. Obviously, $\mathfrak{Q} = \{A \in \mathfrak{U} \mid \bar{\mathfrak{R}}^\sigma A \subset \mathfrak{I}\}$ is a weakly closed two-sided ideal in \mathfrak{U} . Hence, $\mathfrak{Q} = \mathfrak{U}E$ with a central projection $E \in \mathfrak{U}$. From $\mathfrak{R}\mathfrak{I}^*\mathfrak{I} \subset \mathfrak{R}\mathfrak{R}^*\mathfrak{I} \subset \mathfrak{I}$ we obtain $\mathfrak{I}^*\mathfrak{I} \subset \mathfrak{Q}$ and so $\mathfrak{B} \subset \mathfrak{Q}$. On the other hand, $\bar{\mathfrak{R}}^\sigma \mathfrak{Q} \subset \mathfrak{I}$ implies $\mathfrak{I} = \mathfrak{I}\mathfrak{B} \subset \bar{\mathfrak{R}}^\sigma \mathfrak{Q} \subset \mathfrak{I}$. Thus, $\mathfrak{I} = \bar{\mathfrak{R}}^\sigma \mathfrak{Q}$ and therefore $\mathfrak{I} = \bar{\mathfrak{R}}^\sigma \mathfrak{U}E = \bar{\mathfrak{R}}^\sigma E$.

2. THE REPRESENTATION OF A HILBERT MODULE AS A TERNARY RING OF OPERATORS

In the following, we establish two representation theorems which bring Hilbert modules in relation to TROs, providing the essential tool for Sections 3 and 4.

The fundamental results we are going to use can be found in [5]. Most of the basic material is also contained in [8], where Hilbert \mathfrak{U} -modules in the following sense appear as right \mathfrak{U} -rigged spaces:

2.1. DEFINITION. Let \mathfrak{U} be a C^* -algebra. A Hilbert \mathfrak{U} -module $(\mathfrak{H}, \langle \cdot | \cdot \rangle)$ consists of a right \mathfrak{U} -module \mathfrak{H} and a conjugate bilinear map $\langle \cdot | \cdot \rangle: \mathfrak{H} \times \mathfrak{H} \rightarrow \mathfrak{U}$, called the \mathfrak{U} -valued inner product, such that:

- (1) $\langle x | x \rangle \geq 0$ and $\langle x | x \rangle = 0$ only if $x = 0$, $x \in \mathfrak{H}$,
- (2) $\langle x \cdot a | y \rangle = \langle x | y \rangle a$, $x, y \in \mathfrak{H}$, $a \in \mathfrak{U}$,
- (3) $\langle x | y \rangle^* = \langle y | x \rangle$, $x, y \in \mathfrak{H}$,

hold, and such that the norm $\| \cdot \|_{\mathfrak{H}}: x \mapsto \| \langle x | x \rangle \|^{1/2}$ on \mathfrak{H} is complete (cf. [5, Definitions 2.1 and 2.5]). If, in addition, each $\| \cdot \|_{\mathfrak{H}}$ -bounded \mathfrak{U} -module homomorphism of \mathfrak{H} into \mathfrak{U} is of the form $y \mapsto \langle y | x \rangle$ for some $x \in \mathfrak{H}$, then $(\mathfrak{H}, \langle \cdot | \cdot \rangle)$ is called *self-dual over \mathfrak{U}* (cf. [5, paragraph 3]).

For instance, let $\mathfrak{R} \subset \mathcal{L}(H, K)$ be a norm closed TRO and put $\langle X | Y \rangle = Y^*X$. Then $(\mathfrak{R}, \langle \cdot | \cdot \rangle)$ is obviously a Hilbert $[\mathfrak{R}]_H$ -module with norm $\| \cdot \|_{[\mathfrak{R}]_H}$ equal to the operator norm. Moreover, as 4.12 will show, $(\mathfrak{R}, \langle \cdot | \cdot \rangle)$ is self-dual over $[\mathfrak{R}]_H$ if \mathfrak{R} is weakly closed. Conversely, we will prove in Theorem 2.6 that each Hilbert \mathfrak{U} -module $(\mathfrak{H}, \langle \cdot | \cdot \rangle)$ is isomorphic to $(\mathfrak{R}, \langle \cdot | \cdot \rangle)$ for a suitable norm closed TRO \mathfrak{R} . The isomorphism is given by an isometry $U: \mathfrak{H} \rightarrow \mathfrak{R}$ and a $*$ -isomorphism $\pi: \mathfrak{U} \rightarrow [\mathfrak{R}]_H$ satisfying

$$U(x \cdot a) = U(x) \pi(a), \quad U(y)^* U(x) = \pi(\langle x | y \rangle), \quad x, y \in \mathfrak{H}, \quad a \in \mathfrak{U}.$$

In particular, U and π result as the restrictions to \mathfrak{H} and \mathfrak{A} , respectively, of an analogous representation of the self-dual completion \mathfrak{H}'' of \mathfrak{H} over \mathfrak{A}^{**} , which also involves the naturally occurring weak and strong topologies. Then, for a W^* -algebra \mathfrak{A} and a self-dual Hilbert \mathfrak{A} -module $(\mathfrak{H}, \langle \cdot | \cdot \rangle)$, we shall derive in Theorem 2.8 that $(\mathfrak{H}, \langle \cdot | \cdot \rangle)$ is isomorphic to $(\mathfrak{R}, \langle \cdot | \cdot \rangle)$, where \mathfrak{R} is a weakly closed TRO, the isomorphism being likewise a "normal" one. It is easy to see that these theorems contain the well-known representation for C^* - and W^* -algebras by specializing $\mathfrak{H} = \mathfrak{A}$ and $\langle x | y \rangle = y^*x$.

For the remainder we thus fix a C^* -algebra \mathfrak{A} and a Hilbert \mathfrak{A} -module $(\mathfrak{H}, \langle \cdot | \cdot \rangle)$. First, let us recall the self-dual completion \mathfrak{H}'' of \mathfrak{H} .

2.2. PROPOSITION ([5, Corollary 4.3]). *Endowed with the conjugate scalar multiplication $(\lambda, \tau) \mapsto \bar{\lambda}\tau$, the space \mathfrak{H}'' of all $\|\cdot\|_{\mathfrak{H}}$ -bounded \mathfrak{A} -module homomorphisms of \mathfrak{H} into the enveloping W^* -algebra \mathfrak{A}^{**} of \mathfrak{A} becomes a right \mathfrak{A}^{**} -module by*

$$(\tau \cdot A)(x) = A^*\tau(x), \quad \tau \in \mathfrak{H}'', \quad A \in \mathfrak{A}^{**}, \quad x \in \mathfrak{H}.$$

*Moreover, the \mathfrak{A} -valued inner product can be extended to an \mathfrak{A}^{**} -valued inner product on \mathfrak{H}'' , again denoted by $\langle \cdot | \cdot \rangle$, such that:*

- (1) $(\mathfrak{H}'', \langle \cdot | \cdot \rangle)$ is a self-dual Hilbert \mathfrak{A}^{**} -module,
- (2) $\|\cdot\|_{\mathfrak{H}''}$ is equal to the operator norm induced by $\mathcal{L}(\mathfrak{H}, \mathfrak{A}^{**})$,
- (3) $x \mapsto \langle \cdot | x \rangle$ is an isomorphism of $(\mathfrak{H}, \langle \cdot | \cdot \rangle)$ into $(\mathfrak{H}'', \langle \cdot | \cdot \rangle)$,
- (4) $\tau(x) = \langle x | \tau \rangle$, $x \in \mathfrak{H}$, $\tau \in \mathfrak{H}''$,

x and $\langle \cdot | x \rangle$ being identified for each $x \in \mathfrak{H}$.

It is shown in [5, Proposition 3.8] that \mathfrak{H}'' is weak* closed in the dual space $\mathcal{L}(\mathfrak{H}, \mathfrak{A}^{**})$ of the projective tensor product $\mathfrak{H} \hat{\otimes}_{\mathfrak{A}} \mathfrak{A}^*$. Thus, \mathfrak{H}'' is the dual space of a quotient \mathfrak{B} of $\mathfrak{H} \hat{\otimes}_{\mathfrak{A}} \mathfrak{A}^*$ (in the terminology of [7, Corollary 2.13], $\mathfrak{B} = \mathfrak{H} \otimes_{\mathfrak{A}} \mathfrak{A}^*$). As Example 4.12 together with 2.2(1) will prove, the predual of \mathfrak{H}'' is uniquely determined up to isomorphism.

2.3. DEFINITION. Henceforth, we call $\sigma(\mathfrak{H}'', \mathfrak{B})$ the *weak topology* of \mathfrak{H}'' , whereas the *strong topology* of \mathfrak{H}'' will be the topology generated by all seminorms $\tau \mapsto \varphi(\langle \tau | \tau \rangle)^{1/2}$, $0 \leq \varphi \in \mathfrak{A}^*$.

2.4. LEMMA. \mathfrak{H} is strongly dense in \mathfrak{H}'' .

This follows immediately from the arguments in [6, Lemma 2.3].

The proof of Lemma 2.5 starts with simultaneous GNS-constructions for (\mathfrak{A}, φ) and $(\mathfrak{H}, \varphi \circ \langle \cdot | \cdot \rangle)$ given in [5, p. 450].

2.5. LEMMA. Let $0 \leq \varphi \in \mathfrak{A}^*$ and denote the associated cyclic representation of \mathfrak{A}^{**} by $\{\pi_\varphi, \mathfrak{A}_\varphi\}$ (cf. [9, Sect. 1.16]). Then there exist a Hilbert space \mathfrak{H}_φ and a linear map U_φ from \mathfrak{H}'' into $\mathcal{L}(\mathfrak{A}_\varphi, \mathfrak{H}_\varphi)$ with $\|U_\varphi\| \leq 1$ such that:

- (1) $U_\varphi(\tau \cdot A) = U_\varphi(\tau) \pi_\varphi(A)$, $\tau \in \mathfrak{H}''$, $A \in \mathfrak{A}^{**}$,
- (2) $U_\varphi(\eta)^* U_\varphi(\tau) = \pi_\varphi(\langle \tau | \eta \rangle)$, $\tau, \eta \in \mathfrak{H}''$.

Proof. We can suppose that φ is a state. Write N_φ for the kernel $\{x \in \mathfrak{H} \mid \varphi(\langle x | x \rangle) = 0\}$ of the positive Hermitian form $\varphi \circ \langle \cdot | \cdot \rangle$, and denote the quotient map $\mathfrak{H} \rightarrow \mathfrak{H}/N_\varphi$ by $x \mapsto (x)_\varphi$. Then

$$\langle (x)_\varphi | (y)_\varphi \rangle_\varphi = \varphi(\langle x | y \rangle), \quad x, y \in \mathfrak{H}$$

defines a scalar product on \mathfrak{H}/N_φ with corresponding Hilbert space \mathfrak{H}_φ . The same process, applied to \mathfrak{A} and φ , leads to the Hilbert space \mathfrak{A}_φ and the quotient map $a \mapsto (a)_\varphi$.

By the remarks preceeding [5, Proposition 3.1], for each $\tau \in \mathfrak{H}''$ there is a unique $(\tau)_\varphi \in \mathfrak{H}_\varphi$ such that

$$\varphi(\tau(x)) = \langle (x)_\varphi | (\tau)_\varphi \rangle_\varphi, \quad x \in \mathfrak{H}.$$

It follows that for each $\tau \in \mathfrak{H}''$ an operator $U_\varphi(\tau) \in \mathcal{L}(\mathfrak{A}_\varphi, \mathfrak{H}_\varphi)$ with norm $\leq \|\tau\|$ is given by

$$U_\varphi(\tau)(a)_\varphi = (\tau \cdot a)_\varphi, \quad a \in \mathfrak{A}. \quad (2.1)$$

Indeed, [5, Theorem 2.8] yields $\varphi(\tau(x)^* \tau(x)) \leq \|\tau\|^2 \varphi(\langle x | x \rangle)$ for each $x \in \mathfrak{H}$ and therefore,

$$\begin{aligned} \|(\tau \cdot a)_\varphi\|_\varphi &= \sup\{|\langle (x)_\varphi | (\tau \cdot a)_\varphi \rangle_\varphi| \mid \|(x)_\varphi\|_\varphi \leq 1\} \\ &= \sup\{|\varphi(a^* \tau(x))| \mid \|(x)_\varphi\|_\varphi \leq 1\} \\ &\leq \sup\{\varphi(a^* a)^{1/2} \varphi(\tau(x)^* \tau(x))^{1/2} \mid \varphi(\langle x | x \rangle) \leq 1\} \\ &\leq \varphi(a^* a)^{1/2} \|\tau\| = \|(a)_\varphi\|_\varphi \|\tau\|, \end{aligned}$$

so that $U_\varphi(\tau)$ is well defined and continuous on a dense subset of \mathfrak{A}_φ .

Now, (1) results from Eq. (2.1) and the continuity of $U_\varphi(\tau)$, because

$$\begin{aligned} U_\varphi(\tau \cdot A)(a)_\varphi &= ((\tau \cdot A) \cdot a)_\varphi = (\tau \cdot (Aa))_\varphi \\ &= U_\varphi(\tau)(Aa)_\varphi = U_\varphi(\tau) \pi_\varphi(A)(a)_\varphi, \quad a \in \mathfrak{A}. \end{aligned}$$

It remains to show (2). By Lemma 2.4, there is a net $\{x^\alpha\}_\alpha \subset \mathfrak{H}$ which tends to $\tau \in \mathfrak{H}''$ strongly. Since [5, Proposition 2.3] applied to $(\mathfrak{H}'', \langle \cdot | \cdot \rangle)$ gives $\langle \rho | \eta \rangle^* \langle \rho | \eta \rangle \leq \|\eta\|^2 \langle \rho | \rho \rangle$ for all $\rho, \eta \in \mathfrak{H}''$, we obtain

$$\begin{aligned}
\|(\langle \tau \cdot a - x^\alpha \cdot a \mid \eta \rangle)_\omega\|_\omega^2 &= \varphi(\langle \tau \cdot a - x^\alpha \cdot a \mid \eta \rangle^* \langle \tau \cdot a - x^\alpha \cdot a \mid \eta \rangle) \\
&\leq \|\eta\|^2 \varphi(a^* \langle \tau - x^\alpha \mid \tau - x^\alpha \rangle a) \\
&= \|\eta\|^2 \|(\tau \cdot a - x^\alpha \cdot a)_\omega\|_\omega \rightarrow 0.
\end{aligned}$$

Consequently, for each $a \in \mathfrak{A}$

$$\begin{aligned}
\pi_\omega(\langle \tau \mid \eta \rangle)(a)_\omega &= (\langle \tau \cdot a \mid \eta \rangle)_\omega = \lim_\alpha (\langle x^\alpha \cdot a \mid \eta \rangle)_\omega \\
&= \lim_\alpha U_\omega(\eta)^* U_\omega(x^\alpha)(a)_\omega = U_\omega(\eta)^* U_\omega(\tau)(a)_\omega.
\end{aligned}$$

Keeping the notations of Lemma 2.5, we put $\Omega = \{\varphi \in \mathfrak{A}^* \mid 0 \leq \varphi\}$ and define in turn H, K, π, U to be the l^2 -direct sum of $\{\mathfrak{A}_\omega\}_{\omega \in \Omega}$, $\{\mathfrak{H}_\omega\}$, $\{\pi_\omega\}$, $\{U_\omega\}$ (note that $\{\pi, H\}$ is the universal $*$ -representation of \mathfrak{A} extended to \mathfrak{A}^{**} ; cf. [9, Sect. 1.16]). Then we have

2.6. THEOREM. *The map $U: \mathfrak{H}'' \rightarrow \mathcal{L}(H, K)$ is a linear isometry with the following properties:*

- (1) $U(\tau \cdot A) = U(\tau) \pi(A)$, $\tau \in \mathfrak{H}''$, $A \in \mathfrak{A}^{**}$,
- (2) $U(\eta)^* U(\tau) = \pi(\langle \tau \mid \eta \rangle)$, $\tau, \eta \in \mathfrak{H}''$,
- (3) U is homeomorphic in the strong (respectively, weak) topology of \mathfrak{H}'' and the strong (respectively, weak) operator topology;
- (4) $U(\mathfrak{H})$ is a norm closed TRO with weak closure $U(\mathfrak{H}'')$.

Proof. (1) and (2) are clear from 2.5(1), (2) and the definition of U and π . Since π is isometric, (2) implies

$$\|U(\tau)\|^2 = \|\pi(\langle \tau \mid \tau \rangle)\| = \|\tau\|^2, \quad \tau \in \mathfrak{H}''.$$

Next, for each net $\{\tau^\alpha\}_\alpha \subset \mathfrak{H}''$ the following conditions are equivalent:

$$\begin{aligned}
\tau^\alpha &\rightarrow 0 && \text{strongly in } \mathfrak{H}'', \\
\lim_\alpha \varphi(\langle \tau^\alpha \mid \tau^\alpha \rangle) &= 0 && \text{for all } \varphi \in \Omega, \\
\langle \tau^\alpha \mid \tau^\alpha \rangle &\rightarrow 0 && \text{weak } * \text{ in } \mathfrak{A}^{**}, \\
U(\tau^\alpha)^* U(\tau^\alpha) &\rightarrow 0 && \text{weakly in } \mathcal{L}(H), \\
U(\tau^\alpha) &\rightarrow 0 && \text{strongly in } \mathcal{L}(H, K).
\end{aligned}$$

Hence, U is a strong homeomorphism.

Thus, in view of Lemma 2.4, $U(\mathfrak{H}'')$ is contained in the strong closure of $U(\mathfrak{H})$. On the other hand, $U(\mathfrak{H}'')$ is weakly closed. Indeed, suppose X belongs to the weak closure of $U(\mathfrak{H}'')$. Then, $X^*U(\mathfrak{H}'') \subset \pi(\mathfrak{A}^{**})$. Since $\tau \mapsto \pi^{-1}(X^*U(\tau))$ is an \mathfrak{A}^{**} -module homomorphism of \mathfrak{H}'' into \mathfrak{A}^{**} , the self-duality of \mathfrak{H}'' implies the existence of $\eta \in \mathfrak{H}''$ satisfying $\langle \tau | \eta \rangle = \pi^{-1}(X^*U(\tau))$, that is $U(\eta)^*U(\tau) = X^*U(\tau)$, $\tau \in \mathfrak{H}''$. Hence, $X = U(\eta) \in U(\mathfrak{H}'')$. Therefore, $U(\mathfrak{H}'')$ is the weak-operator closure of $U(\mathfrak{H})$.

It remains to show that U is a weak homeomorphism. To this end we require two lemmas.

LEMMA 2.6.1. U^{-1} is weakly continuous.

Proof. Fix $f \in \mathfrak{B}$ and write $f = f_1 + f_2 + i(f_3 + f_4)$, each f_k being of the form $f_k = \sum_{n \in \mathbb{N}} x_{n,k} \otimes \varphi_{n,k}$ with certain sequences $\{x_{n,k}\}_n \subset \mathfrak{H}$, $\{\varphi_{n,k}\}_n \subset \Omega$ satisfying $\sum_n \|x_{n,k}\|_{\mathfrak{H}} \|\varphi_{n,k}\| < \infty$. Now, fix k and define $y_n = x_{n,k}$, $\psi_n = \varphi_{n,k}$, $n \in \mathbb{N}$. Then the functional $X \mapsto \sum_{n \in \mathbb{N}} \langle X(I)_{\psi_n} | (y_n)_{\psi_n} \rangle_{\psi_n}$ is weak-operator continuous, because

$$\sum_{n \geq N} |\langle X(I)_{\psi_n} | (y_n)_{\psi_n} \rangle_{\psi_n}| \leq \|X\| \sum_{n \geq N} \|\psi_n\| \|y_n\|_{\mathfrak{H}}$$

and the latter tends to zero as $N \rightarrow \infty$. Since for each $\tau \in \mathfrak{H}''$

$$\sum_n \langle U(\tau)(I)_{\psi_n} | (y_n)_{\psi_n} \rangle_{\psi_n} = \sum_n \psi_n(\langle \tau | y_n \rangle) = f_k(\tau)$$

holds, the assertion follows.

LEMMA 2.6.2. A linear functional f on \mathfrak{H}'' is weakly continuous if, and only if, it is strongly continuous on $D_1(\mathfrak{H}'')$.

Proof. First, let $A \subset \mathfrak{H}''$ be convex and strongly closed. Since U is a strong homeomorphism, $U(A)$ is strongly, and hence weakly, closed in $\mathcal{L}(H, K)$. By Lemma 2.6.1, A is weakly closed. Next, suppose f is strongly continuous on $D_1(\mathfrak{H}'')$. Then $\{\tau \in D_1(\mathfrak{H}'') | f(\tau) = 0\}$ is convex and weakly closed and so, f is weakly continuous.

The weak continuity of U now follows from 2.6.2 and the observation that U is weakly continuous on norm bounded sets, which in turn is immediate from Eq. (2.1) and the definition of \mathfrak{B} :

$$\begin{aligned} \langle (x)_{\omega} | U(\tau)(a)_{\omega} \rangle_K &= \langle (x)_{\omega} | U_{\omega}(\tau)(a)_{\omega} \rangle_{\omega} = \varphi(a^* \tau(x)), \\ x \in \mathfrak{B}, \quad a \in \mathfrak{A}, \quad \tau \in \mathfrak{H}'', \quad \text{and} \quad \varphi \in \Omega. \end{aligned}$$

2.7. COROLLARY. The unit ball of \mathfrak{H}'' is both weakly and strongly dense in the unit ball of \mathfrak{H}'' .

Proof. By 1.4, $D_1(U(\mathfrak{H}))$ is both weakly and strongly dense in $D_1(U(\mathfrak{H}''))$. The assertion now follows from 2.6(3).

We conclude this section with the analogue of Theorem 2.6 for self-dual Hilbert modules over W^* -algebras. Since the proof is very similar to that of 2.6, we omit the details.

2.8. THEOREM. *Suppose \mathfrak{A} is a W^* -algebra and $(\mathfrak{H}, \langle \cdot | \cdot \rangle)$ is a self-dual Hilbert \mathfrak{A} -module. Put $\Lambda = \{\varphi \in \mathfrak{A}_* \mid 0 \leq \varphi\}$ and, in the notations of Lemma 2.5, define H, K, π, U to be the l^2 -direct sum of $\{\mathfrak{A}_\varphi\}_{\varphi \in \Lambda}$, $\{\mathfrak{H}_\varphi\}$, $\{\pi_\varphi| \mathfrak{A}\}$, $\{U_\varphi| \mathfrak{H}\}$, respectively. Then $U: \mathfrak{H} \rightarrow \mathcal{L}(H, K)$ is a linear isometry, and*

$$(1) \quad U(x \cdot a) = U(x) \pi(a), \quad x \in \mathfrak{H}, a \in \mathfrak{A};$$

$$(2) \quad U(y)^* U(x) = \pi(\langle x | y \rangle), \quad x, y \in \mathfrak{H};$$

(3) *U is a homeomorphism in the topology generated by all seminorms $x \mapsto \varphi(\langle x | x \rangle)^{1/2}$, $\varphi \in \Lambda$ (the topology generated by $\mathfrak{H} \hat{\otimes} \mathfrak{A}_*$, respectively) and the strong-operator topology (the weak-operator topology, respectively);*

$$(4) \quad U(\mathfrak{H}) \text{ is a weakly closed TRO.}$$

3. THE REPRESENTATION THEOREM FOR TERNARY C^* -RINGS

A ternary C^* -ring $(\mathfrak{X}, (\cdot, \cdot, \cdot), \|\cdot\|)$ will be called *isomorphic* (respectively, *anti-isomorphic*) to a TRO, if there are Hilbert spaces H and K and a linear isometry $U: \mathfrak{X} \rightarrow \mathcal{L}(H, K)$ satisfying

$$U((u, y, z)) = U(x) U(y)^* U(z), \quad x, y, z \in \mathfrak{X}$$

(respectively, $U((x, y, z)) = -U(x) U(y)^* U(z)$, $x, y, z \in \mathfrak{X}$). Obviously, $U(\mathfrak{X})$ is a norm closed TRO. The purpose of this section is now the proof of

3.1. REPRESENTATION THEOREM. *Let $(\mathfrak{X}, (\cdot, \cdot, \cdot), \|\cdot\|)$ be a ternary C^* -ring.*

(1) *Then \mathfrak{X} is the direct sum of two ternary C^* -subrings \mathfrak{X}_+ and \mathfrak{X}_- in such a way that \mathfrak{X}_+ (respectively, \mathfrak{X}_-) is isomorphic (respectively, anti-isomorphic) to a TRO. Moreover, this decomposition of \mathfrak{X} is unique.*

(2) *If P denotes the projection of \mathfrak{X} onto \mathfrak{X}_+ with kernel \mathfrak{X}_- , then $2P - 1$ is the only operator $T \in \mathcal{L}(\mathfrak{X})$ which satisfies $T^2 = 1$ and $T((x, y, z)) = (Tx, y, z) = (x, Ty, z) = (x, y, Tz)$, $x, y, z \in \mathfrak{X}$ and which has the property that $(\mathfrak{X}, T \circ (\cdot, \cdot, \cdot), \|\cdot\|)$ is isomorphic to a TRO.*

The proof is carried out in several steps (3.2–3.7). The first one shows that a ternary C^* -ring is a special module over a C^* -algebra:

3.2. PROPOSITION. Let $(\mathfrak{X}, (\cdot, \cdot, \cdot), \|\cdot\|)$ be a ternary C^* -ring. Then there exists a unique pair $(\mathfrak{A}, \mathfrak{a})$ such that

(1) \mathfrak{A} is a C^* -algebra, and \mathfrak{X} is a right Banach \mathfrak{A} -module,

(2) $\mathfrak{a}: \mathfrak{X} \times \mathfrak{X} \rightarrow \mathfrak{A}$ is conjugate bilinear with $\|\mathfrak{a}\| \leq 1$ and

$$\mathfrak{a}(x \cdot a, y) = \mathfrak{a}(x, y)a, \quad x, y \in \mathfrak{X}, \quad a \in \mathfrak{A},$$

$$\mathfrak{a}(x, y)^* = \mathfrak{a}(y, x), \quad x, y \in \mathfrak{X},$$

(3) $(x, y, z) = x \cdot \mathfrak{a}(z, y)$, $x, y, z \in \mathfrak{X}$,

(4) the linear hull of $\mathfrak{a}(\mathfrak{X}, \mathfrak{X})$ is norm dense in \mathfrak{A} .

Moreover, we have $\|x\|^2 = \|\mathfrak{a}(x, x)\|$ for each $x \in \mathfrak{X}$.

Proof. The last statement is a consequence of Eq. (0.5), condition 3.2(3), and the fact that $\|\mathfrak{a}\| \leq 1$:

$$\|x\|^3 = \|x \cdot \mathfrak{a}(x, x)\| \leq \|x\| \|\mathfrak{a}(x, x)\| \leq \|x\|^3, \quad x \in \mathfrak{X}.$$

Existence. To begin with, let V denote the linear hull of $\{(\cdot, x, y) \mid x, y \in \mathfrak{X}\}$ in $\mathcal{L}(\mathfrak{X})$; observe that $(\cdot, x, y) \in \mathcal{L}(\mathfrak{X})$ by (0.1) and (0.4). We show that V is a pre- C^* -algebra in the operator norm.

From (0.3) it is apparent that

$$(\cdot, x_2, y_2) \circ (\cdot, x_1, y_1) = (\cdot, (x_2, y_1, x_1), y_2)$$

and thus, V is a subalgebra of $\mathcal{L}(\mathfrak{X})$. Furthermore,

$$j \left(\sum_{i=1}^n (\cdot, x_i, y_i) \right) = \sum_{i=1}^n (\cdot, y_i, x_i)$$

defines an involution j on V . In fact, if $\sum_i (\cdot, x_i, y_i)$ vanishes on \mathfrak{X} , set $A = \sum_i (\cdot, y_i, x_i)$ and use (0.2), (0.3) to obtain

$$\begin{aligned} (Az, Az, Az) &= \sum_{i,j,k} ((z, y_i, x_i), (z, y_j, x_j), (z, y_k, x_k)) \\ &= \sum_{i,j,k} (z, ((z, y_j, x_j), x_i, y_i), (z, y_k, x_k)) \\ &= \sum_{j,k} \left(z, \sum_i ((z, y_j, x_j), x_i, y_i), (z, y_k, x_k) \right) = 0, \quad z \in \mathfrak{X} \end{aligned}$$

so that j is well defined. Clearly, j is conjugate linear and satisfies $j^2 = 1$, while with $A_i = (\cdot, x_i, y_i)$, $i = 1, 2$ we have

$$\begin{aligned} j(A_2 \circ A_1)z &= j((\cdot, (x_2, y_1, x_1), y_2))z = (z, y_2, (x_2, y_1, x_1)) \\ &= ((z, y_2, x_2), y_1, x_1) = j(A_1) \circ j(A_2)z, \quad z \in \mathfrak{X} \end{aligned}$$

and so, j is anti-multiplicative. Finally, relative to " \circ " and j , the norm induced by $\mathcal{L}(\mathfrak{X})$ on V has the C^* -property, for, if we put $A = \sum_{i=1}^n (\cdot, x_i, y_i)$, then, by (0.5) and (0.4),

$$\begin{aligned} \|Az\|^3 &= \|(Az, Az, Az)\| = \left\| \sum_i ((z, x_i, y_i), Az, Az) \right\| \\ &= \left\| \sum_i (z, (Az, y_i, x_i), Az) \right\| = \|(z, j(A) \circ Az, Az)\| \\ &\leq \|z\| \|j(A) \circ Az\| \|Az\| \leq \|z\|^2 \|j(A) \circ A\| \|Az\|, \quad z \in \mathfrak{X}, \end{aligned}$$

which implies that $\|A\|^2 \leq \|j(A) \circ A\|$.

Next, define \mathfrak{A} to be the opposite algebra of the norm closure of V in $\mathcal{L}(\mathfrak{X})$ and denote the multiplication and involution on \mathfrak{A} by $(a, b) \mapsto ab$ and $a \mapsto a^*$, respectively. Then \mathfrak{A} is a C^* -algebra, and \mathfrak{X} becomes a faithful right Banach \mathfrak{A} -module by $x \cdot a = a(x)$. If we now introduce $\alpha: \mathfrak{X} \times \mathfrak{X} \mapsto \mathfrak{A}$ according to

$$\alpha(z, y) = (\cdot, y, z), \quad y, z \in \mathfrak{X}, \quad (3.1)$$

then (3) and (4) are clear by definition, while, obviously, α is conjugate linear with $\|\alpha\| \leq 1$ and $\alpha(x, y)^* = \alpha(y, x)$. Moreover, writing $a = \alpha(x_1, y_1)$, we conclude that for all $x, y, z \in \mathfrak{X}$

$$x \cdot \alpha(y \cdot a, z) = (x, z, (y, y_1, x_1)) = ((x, z, y), y_1, x_1) = x \cdot \alpha(y, z)a,$$

hence, $\alpha(y \cdot a, z) = \alpha(y, z)a$. It follows that $\alpha(y \cdot a, z) = \alpha(y, z)a$ for all $y, z \in \mathfrak{X}$ and $a \in \mathfrak{A}$.

Uniqueness. Suppose, (\mathfrak{B}, b) is another pair satisfying (1)–(4) for \mathfrak{X} . First, the module action of \mathfrak{B} on \mathfrak{X} gives rise to an anti-homomorphism $m: \mathfrak{B} \rightarrow \mathcal{L}(\mathfrak{X})$. By (3) and (4), $m(b(x, y)) = \alpha(x, y)$. Furthermore, m is a $*$ -homomorphism, because $m(b(x, y)^*) = \alpha(y, x) = \alpha(x, y)^* = m(b(x, y))^*$. Finally,

$$m(\mathfrak{B}) = \overline{m(\lim b(\mathfrak{X}, \mathfrak{X})^n)} \subset \overline{\lim m(b(\mathfrak{X}, \mathfrak{X}))^n} = \mathfrak{A}.$$

Thus, m is a $*$ -isomorphism of \mathfrak{B} onto \mathfrak{A} which transforms b into α .

With the notations of 3.2, we are now going to show that

$$\mathfrak{X}_+ = \{x \in \mathfrak{X} \mid \alpha(x, x) \geq 0\} \quad \text{and} \quad \mathfrak{X}_- = \{x \in \mathfrak{X} \mid \alpha(x, x) \leq 0\}$$

define two α -orthogonal \mathfrak{A} -submodules which decompose \mathfrak{X} . Besides orthogonality the crucial point consists in proving that \mathfrak{X}_+ and \mathfrak{X}_- are linear subspaces of \mathfrak{X} . To this end we consider not only the "right" structure of (\cdot, \cdot, \cdot) , as we did in Proposition 3.2, but also the "left" one which is

provided by the C^* -algebra of all α -adjoinable operators on \mathfrak{X} and another form b (cf. Proposition 3.4). For positive α , this C^* -algebra has already been studied in [5, 6, 8].

3.3. DEFINITION. Suppose F and G are two maps on \mathfrak{X} satisfying $\alpha(Fx, y) = \alpha(x, Gy)$ for all $x, y \in \mathfrak{X}$. By 3.2, both F and G are continuous \mathfrak{U} -module homomorphisms which determine each other uniquely. Hence, we call G the α -adjoint F^* of F .

By $\mathfrak{U}_\alpha(\mathfrak{X})$ we shall denote the $*$ -algebra of all maps F on \mathfrak{X} which possess an α -adjoint. Using 3.2 again, it follows immediately that $\mathfrak{U}_\alpha(\mathfrak{X})$ is a C^* -algebra with unit $I = I_{\mathfrak{X}}$ in the norm inherited from $\mathcal{L}(\mathfrak{X})$. Observe that \mathfrak{X} is a left Banach $\mathfrak{U}_\alpha(\mathfrak{X})$ -module.

3.4. PROPOSITION. Define $b: \mathfrak{X} \times \mathfrak{X} \rightarrow \mathfrak{U}_\alpha(\mathfrak{X})$ by $b(x, y) \cdot z = x \cdot \alpha(z, y)$, $x, y \in \mathfrak{X}$ (that is, $b(x, y) = (x, y, \cdot)$). Then

(1) b is conjugate bilinear with $\|b\| \leq 1$ and $b(F \cdot x, y) = Fb(x, y)$, $b(x, y)^* = b(y, x)$, $\|x\|^2 = \|b(x, x)\|$, $x, y \in \mathfrak{X}$, $F \in \mathfrak{U}_\alpha(\mathfrak{X})$.

(2) If \mathfrak{B} denotes the C^* -algebra generated by $b(\mathfrak{X}, \mathfrak{X})$, then $\mathfrak{U}_\alpha(\mathfrak{X})$ is the double centralizer algebra $M(\mathfrak{B})$ of \mathfrak{B} .

Further, let $\mathfrak{U}_b(\mathfrak{X})$ denote the C^* -algebra of all maps on \mathfrak{X} which possess a b -adjoint. Then

(3) $\mathfrak{U}_b(\mathfrak{X})$ is the double centralizer algebra of the opposite algebra \mathfrak{U}_\sim of \mathfrak{U} , and the form $c: \mathfrak{X} \times \mathfrak{X} \rightarrow \mathfrak{U}_b(\mathfrak{X})$ constructed for b in analogy with the process $\alpha \rightsquigarrow b$ coincides with a .

(4) $M(\mathfrak{B})$ commutes with $M(\mathfrak{U}_\sim)$.

Proof. Since, by Eq. (3.1),

$$\begin{aligned} \alpha(b(x, y) \cdot z_1, z_2) &= \alpha((x, y, z_1), z_2) = (\cdot, z_2, (x, y, z_1)) \\ &= (\cdot, (y, x, z_2), z_1) = \alpha(z_1, b(y, x) \cdot z_2), \quad x, y, z_1, z_2 \in \mathfrak{X}, \end{aligned}$$

it follows that $b(x, y) \in \mathfrak{U}_\alpha(\mathfrak{X})$ and $b(x, y)^* = b(y, x)$. For each continuous \mathfrak{U} -module homomorphism $F: \mathfrak{X} \rightarrow \mathfrak{X}$ we obtain

$$b(F(x), y) \cdot z = F(x) \cdot \alpha(z, y) = F(x \cdot \alpha(z, y)) = F(b(x, y) \cdot z), \quad x, y, z \in \mathfrak{X}.$$

In particular, \mathfrak{B} is a two-sided ideal in $\mathfrak{U}_\alpha(\mathfrak{X})$.

As (1) is clear, we turn to the form c for the system $(\mathfrak{X}, \mathfrak{B}, b)$. From $c(x, y)(z) = b(z, y) \cdot x = z \cdot \alpha(x, y) = \alpha(x, y)(z)$ we conclude that $c = \alpha$. Thus, the C^* -algebra generated by $c(\mathfrak{X}, \mathfrak{X})$ is the anti-homomorphic image \mathfrak{U}_\sim of \mathfrak{U} in $\mathcal{L}(\mathfrak{X})$. As above, \mathfrak{U}_\sim is a two-sided ideal in $\mathfrak{U}_b(\mathfrak{X})$. Now, put $\mathfrak{U}_b = \mathfrak{U}_b(\mathfrak{X})$. In view of [1, Proposition 3.7], we may identify $M(\mathfrak{U}_\sim)$ with \mathfrak{U}_b .

if the annihilator of \mathfrak{A}_\sim in \mathfrak{A}_b is trivial and \mathfrak{A}_b is complete in the strict \mathfrak{A}_\sim -topology.

So, fix $F \in \mathfrak{A}_b$ with $F\mathfrak{A}_\sim = \{0\}$. Then, $F(x \cdot a) = 0$ for all $x \in \mathfrak{X}$ and $a \in \mathfrak{A}$. Since, by Cohen's factorization theorem [13, p. 80], we have $\mathfrak{X} = \mathfrak{X} \cdot \mathfrak{A}$ it follows that $F = 0$. Next, let $\{F_\alpha\}_\alpha \subset \mathfrak{A}_b$ be a Cauchy net in the strict \mathfrak{A}_\sim -topology, which means that

$$0 = \lim_{\alpha, \alpha'} \|(F_\alpha - F_{\alpha'}) \cdot a\| = \lim_{\alpha, \alpha'} \|a \cdot (F_\alpha - F_{\alpha'})\|, \quad a \in \mathfrak{A}.$$

Using Cohen's theorem again we obtain $0 = \lim_{\alpha, \alpha'} \|(F_\alpha - F_{\alpha'})(x)\|$, $x \in \mathfrak{X}$. Similarly, $\lim_\alpha F_\alpha^*(x)$ exists for all $x \in \mathfrak{X}$, because

$$\begin{aligned} \|(F_\alpha^* - F_{\alpha'}^*)(x \cdot a)\| &= \sup\{\|b((F_\alpha^* - F_{\alpha'}^*)(x \cdot a), y)\| \mid y \in D_1(\mathfrak{X})\} \\ &= \sup\{\|b(x \cdot a, (F_\alpha - F_{\alpha'})(y))\| \mid y \in D_1(\mathfrak{X})\} \\ &\leq \|x\| \sup\{\|(F_\alpha - F_{\alpha'})(y) \cdot a^*\| \mid y \in D_1(\mathfrak{X})\} \\ &= \|x\| \|a^*(F_\alpha - F_{\alpha'})\| \xrightarrow{\alpha, \alpha'} 0. \end{aligned}$$

Now, the operator $F: \mathfrak{X} \rightarrow \mathfrak{X}$, defined by $F(x) = \lim_\alpha F_\alpha(x)$ is linear, while $b(F(x), y) = \lim_\alpha b(F_\alpha(x), y) = b(x, \lim_\alpha F_\alpha^*(y))$ yields $F \in \mathfrak{A}_b$. Finally, $\{F_\alpha\}_\alpha$ tends strictly to F , because

$$F \cdot a(x) = F(x \cdot a) = \lim_\alpha F_\alpha(x \cdot a) = \left(\lim_\alpha F_\alpha \cdot a \right)(x), \quad x \in \mathfrak{X}, a \in \mathfrak{A}.$$

The proof of $M(\mathfrak{B}) = \mathfrak{A}_a(\mathfrak{X})$ is similar. It remains to show (4). So, take $F = F^* \in \mathfrak{A}_a(\mathfrak{X})$. Then $FG = GF$ follows from

$$\begin{aligned} z \cdot a(FG(x), y) &= z \cdot a(G(x), F(y)) = b(z, F(y)) \cdot G(x) \\ &= G(b(z, F(y)) \cdot x) = G(z \cdot a(F(x), y)) = G(b(z, y) \cdot F(x)) \\ &= b(z, y) \cdot GF(x) = z \cdot a(GF(x), y). \end{aligned}$$

3.5. LEMMA. *Let $x \in \mathfrak{X}$, and let \mathfrak{T} denote the ternary C^* -ring generated by x . There exist $u, v \in \mathfrak{T}$ such that $x = u + v$, $a(u, v) = 0$, $a(u, u) = a(x, x)_+$ and $a(v, v) = -a(x, x)_-$.*

Proof. We may assume that $\|x\| \leq 1$. Put $a = a(x, x)$. Then $x_n = x \cdot a_+^{1/n}$ is in \mathfrak{T} for all $n \in \mathbb{N}$. Moreover,

$$\|x_n - x_m\|^2 = \|a(x_n - x_m, x_n - x_m)\| = \|a_+^{1/2}(a_+^{1/n} - a_+^{1/m})\|^2.$$

Now let \mathfrak{C} denote the C^* -algebra generated by a . By Dini's theorem, $\{a_+^{1/2+1/n}\}_n$ tends uniformly to $a_+^{1/2}$ in \mathfrak{C} . Hence, the norm limit

$u = \lim_n x_n \in \mathfrak{I}$ exists and satisfies $\alpha(u, u) = \lim_n \alpha(x_n, x_n) = a_+$. Similarly, $v = \lim_n x \cdot a^{1/n} \in \mathfrak{I}$ and $\alpha(v, v) = -a_-$. From $a_+ a_- = 0$ we conclude that $\alpha(u, v) = 0$. Finally, representing $\mathfrak{C} \oplus \mathbb{C}$ as the continuous functions on $[-1, 1]$, we easily obtain that $\{e_n\}_n = \{a_+^{1/n} + a_-^{1/n}\}_n$ is an approximate identity for \mathfrak{C} . This gives us

$$\alpha(y, u + v) = \alpha(y, \lim_n x \cdot e_n) = \lim_n \alpha(y, x) e_n = \alpha(y, x), \quad y \in \mathfrak{I},$$

and hence $x = u + v$.

3.6. LEMMA. *For each $x \in \mathfrak{X}_\pm$ there exists $x' \in \mathfrak{X}_\pm$ satisfying $\alpha(x', x') = \pm(\pm\alpha(x, x))^{1/2}$ and $x = x' \cdot (\pm\alpha(x, x))^{1/4}$.*

Proof. It suffices to consider the case $x \in \mathfrak{X}_+$. First, we have

LEMMA 3.6.1. *The ternary C^* -ring \mathfrak{I} generated by x is contained in \mathfrak{X}_+ , and $(\mathfrak{I}, \alpha | \mathfrak{I} \times \mathfrak{I})$ is a Hilbert \mathfrak{C} -module, where \mathfrak{C} denotes the C^* -algebra generated by $a = \alpha(x, x)$.*

Proof. Indeed, define inductively $x_1 = x$, $x^{2i+1} = (x^{2i} - 1, x, x)$, $i \in \mathbb{N}$, and let $y = \sum_{i=0}^n \lambda_i x^{2i+1}$ be a ternary polynomial in x ; then

$$\begin{aligned} \alpha(y, y) &= \sum_{i,j} \lambda_i \bar{\lambda}_j \alpha(x^{2i+1}, x^{2j+1}) = \sum_{i,j} \lambda_i \bar{\lambda}_j \alpha(x \cdot a^i, x \cdot a^j) \\ &= \sum_{i,j} \lambda_i \bar{\lambda}_j a^i a^j = \left(\sum_{i=0}^n \lambda_i a^i \right) \left(\sum_{i=0}^n \lambda_i a^i \right)^* a \geq 0 \end{aligned}$$

by 3.2(2), hence, $\alpha(y, y) \geq 0$ for all $y \in \mathfrak{I}$ so that 3.6.1 follows.

Applying Theorem 2.6 to the Hilbert \mathfrak{C} -module \mathfrak{I} in 3.6.1, we find Hilbert spaces H and K , a linear isometry $U: \mathfrak{I} \rightarrow \mathscr{L}(H, K)$ and a faithful $*$ -representation $\{\pi, H\}$ of \mathfrak{C} such that

$$U(y \cdot a) = U(y) \pi(a), \quad U(z)^* U(y) = \pi(\alpha(y, z)), \quad y, z \in \mathfrak{I}, \quad a \in \mathfrak{C}.$$

Now, let R be the partial isometry of the polar decomposition of $X = U(x)$. By Notations 1.1 and Proposition 1.2, $Y \mapsto R^* Y$ is an isomorphism of $U(\mathfrak{I})$ onto $\pi(\mathfrak{C})$. Defining $x' = U^{-1}(R |X|^{1/2})$ we obtain

$$\alpha(x', x') = \pi^{-1}(U(x')^* U(x')) = \pi^{-1}(|X|) = \alpha(x, x)^{1/2}$$

and

$$x' \cdot \alpha(x, x)^{1/4} = U^{-1}(R |X|^{1/2} \pi(\alpha(x, x))^{1/4}) = U^{-1}(X) = x.$$

3.7. PROPOSITION. \mathfrak{X}_+ and \mathfrak{X}_- are ternary C^* -subrings of \mathfrak{X} , and $\mathfrak{X} = \mathfrak{X}_+ \oplus \mathfrak{X}_-$. The associated projection $P: \mathfrak{X} \rightarrow \mathfrak{X}_+$ is self-adjoint relative to a and b .

Proof. We first show that

$$\mathfrak{X}_{\pm} = \{x \in \mathfrak{X} \mid b(x, x) \geq 0\}. \quad (3.2)$$

So, let $x \in \mathfrak{X}_+$ and choose $x' \in \mathfrak{X}_+$ according to Lemma 3.6. Then

$$\begin{aligned} b(x', x')^2 &= b(b(x', x') \cdot x', x') = b(x' \cdot a(x', x'), x') \\ &= b(x' \cdot a(x, x)^{1/2}, x') \\ &= b(x' \cdot a(x, x)^{1/4}, x' \cdot a(x, x)^{1/4}) = b(x, x), \end{aligned}$$

and thus, $b(x, x) \geq 0$. Similarly, $b(x, x) \leq 0$ for each $x \in \mathfrak{X}_-$. Now, Eq. (3.2) follows from the identity $a = c$ in 3.4(3) and the fact that the positive cones of \mathfrak{A}_- and \mathfrak{A} coincide.

Let $x \in \mathfrak{X}_+$, $y \in \mathfrak{X}_-$. Then $y \cdot a \in \mathfrak{X}_-$, where $a = a(x, x)^{1/2}$. By 3.4(1), $0 \geq b(y \cdot a, y \cdot a) = b(y \cdot a(x, x), y) = b(b(y, x) \cdot x, y) = b(x, y) * b(x, y)$ and hence, $b(x, y) = 0$. Therefore, $b(\mathfrak{X}_+, \mathfrak{X}_-) = \{0\}$.

Next, let $x, y \in \mathfrak{X}_+$. By Lemma 3.5, we have $x + y = u + v$ with $u \in \mathfrak{X}_+$, $v \in \mathfrak{X}_-$. Since $a(\mathfrak{X}_+, \mathfrak{X}_-) = \{0\}$ we obtain $0 = a(x, v) = a(y, v) = a(u, v)$. Hence, $a(v, v) = a(x + y - u, v) = 0$ and so, $x + y = u \in \mathfrak{X}_+$. Moreover, using $a(x \cdot a, x \cdot a) = a * a(x, x)a \geq 0$, $x \in \mathfrak{X}_+$, $a \in \mathfrak{A}$, we conclude that

$$a((x, y, z), (x, y, z)) = a(x \cdot a(z, y), x \cdot a(z, y)) \geq 0, \quad x, y, z \in \mathfrak{X}_+.$$

Thus, \mathfrak{X}_+ is a ternary subring of \mathfrak{X} which is norm closed, because the positive cone of \mathfrak{A} is closed and a is continuous. Similarly, \mathfrak{X}_- is a ternary C^* -subring of \mathfrak{X} . Finally, since \mathfrak{X}_+ and \mathfrak{X}_- are a - and b -orthogonal the projection P is a - and b -self-adjoint.

Proof of 3.1. We start with 3.1(1). Define $\alpha_+ = a|_{\mathfrak{X}_+ \times \mathfrak{X}_+}$ and let \mathfrak{A}_1 denote the C^* -algebra generated by $\alpha_+(\mathfrak{X}_+, \mathfrak{X}_+)$. By Proposition 3.7, $(\mathfrak{X}_+, \alpha_+)$ is a faithful Hilbert \mathfrak{A}_1 -module. In view of Theorem 2.6, there exist Hilbert spaces H_+ and K_+ and an isometry $U_+ : \mathfrak{X}_+ \rightarrow \mathcal{L}(H_+, K_+)$ such that $\alpha_+(x, y) \mapsto U_+(y) * U_+(x)$ provides a faithful $*$ -representation of \mathfrak{A}_1 on H_+ and $U_+((x, y, z)) = U_+(x \cdot \alpha_+(z, y)) = U_+(x) U_+(y) * U_+(z)$. Hence, \mathfrak{X}_+ is isomorphic to a TRO. By considering $-\alpha|_{\mathfrak{X}_- \times \mathfrak{X}_-}$ we see that \mathfrak{X}_- is anti-isomorphic to a TRO.

To show uniqueness, assume \mathfrak{X}_1 and \mathfrak{X}_2 are ternary C^* -subrings of \mathfrak{X} with $\mathfrak{X} = \mathfrak{X}_1 \oplus \mathfrak{X}_2$ such that representations $U_i: \mathfrak{X}_i \rightarrow \mathcal{L}(H_i, K_i)$ $i = 1, 2$, of the

prescribed kind exist. Put $\mathfrak{A}_1 = [U_1(\mathfrak{X}_1)]_{H_1}$. Then \mathfrak{X}_1 becomes a right Banach \mathfrak{A}_1 -module by

$$x \cdot A = U_1^{-1}(U_1(x)A), \quad x \in \mathfrak{X}_1, \quad A \in \mathfrak{A}_1.$$

Define a continuous Hermitian form $a_1: \mathfrak{X}_1 \times \mathfrak{X}_1 \rightarrow \mathfrak{A}_1$ according to

$$a_1(x, y) = U_1(y)^* U_1(x), \quad x, y \in \mathfrak{X}_1.$$

Since $x \cdot a_1(z, y) = U_1^{-1}(U_1(x) U_1(y)^* U_1(z)) = (x, y, z)$, the uniqueness argument of Proposition 3.2 implies that $a(x, y) \mapsto a_1(x, y)$ defines a $*$ -isomorphism of the C^* -algebra generated by $a(\mathfrak{X}_1, \mathfrak{X}_1)$ onto \mathfrak{A}_1 . Because of $a_1(x, x) \geq 0$ we obtain $\mathfrak{X}_1 \subset \mathfrak{X}_+$. Similarly, $\mathfrak{X}_2 \subset \mathfrak{X}_-$. Thus, $\mathfrak{X}_+ = \mathfrak{X}_1$ and $\mathfrak{X}_- = \mathfrak{X}_2$.

To establish 3.1(2), put $T = 2P - I$. Since, by 3.7, T is a - (respectively, b -) self-adjoint it is easy to check that T has the required properties. Moreover, $a_T: (x, y) \mapsto a(Tx, y)$ is an \mathfrak{A} -valued inner product on \mathfrak{X} , because

$$a_T(x, x) = a(2Px - x, x) = 2a(x, x)_+ - a(x, x) = |a(x, x)| \geq 0, \quad x \in \mathfrak{X}.$$

Using Eq. (2.6) with $(\mathfrak{H}, \langle \cdot | \cdot \rangle) = (\mathfrak{X}, a_T)$, we get Hilbert spaces H and K and an isometry $U: \mathfrak{X} \rightarrow \mathcal{L}(H, K)$ such that $U(x) U(y)^* U(z) = U(x \cdot a_T(z, y)) = U((x, y), Tz) = U \circ T((x, y, z))$.

Finally, suppose that $S: \mathfrak{X} \rightarrow \mathfrak{X}$ and $V: \mathfrak{X} \rightarrow \mathcal{L}(H_S, K_S)$ have the same properties as T and U . Obviously, $S \in \mathfrak{A}_a(\mathfrak{X})$ is self-adjoint. Hence, $S = E - F$, $1 = E + F$, $EF = 0$ with self-adjoint idempotents $E, F \in \mathfrak{A}_a(\mathfrak{X})$. In particular, E and F are \mathfrak{A} -module homomorphisms. Thus, $\mathfrak{X}_1 = E(\mathfrak{X})$ and $\mathfrak{X}_2 = F(\mathfrak{X})$ are ternary C^* -subrings of \mathfrak{X} , and $\mathfrak{X} = \mathfrak{X}_1 \oplus \mathfrak{X}_2$. Furthermore, $V_1 = V|_{\mathfrak{X}_1}$ is an isomorphism. Because of $V((x, y, z)) = -V \circ S(F(x, y, z)) = -V(x) V(y)^* V(z)$ for all $x, y, z \in \mathfrak{X}_2$, $V_2 = V|_{\mathfrak{X}_2}$ is an anti-isomorphism. Now, 3.1(1) implies $\mathfrak{X}_+ = \mathfrak{X}_1$, $\mathfrak{X}_- = \mathfrak{X}_2$ and $S = P - (I - P) = T$.

To give examples, first let \mathfrak{R} be a norm closed TRO between H and K . Then $(\mathfrak{R}, (\cdot, \cdot, \cdot), \|\cdot\|)$ is a ternary C^* -ring, where $\|\cdot\|$ denotes the operator norm and (\cdot, \cdot, \cdot) is given by $(A, B, C) = AB^*C$. In particular, this way each C^* -algebra is a ternary C^* -ring. If $(\mathfrak{H}, \langle \cdot | \cdot \rangle)$ is a Hilbert \mathfrak{A} -module (or, more specifically, a complex Hilbert space), then $(\mathfrak{H}, (\cdot, \cdot, \cdot), \|\cdot\|_{\mathfrak{H}})$ is a ternary C^* -ring with $(x, y, z) = x \cdot \langle z | y \rangle$. In all these situations we obtain $T = I$ if theorem 3.1 is applied. It is, however, easy to see that T and I can be different.

3.8. EXAMPLE. Suppose Ω is a compact Hausdorff space which contains a nonvoid open and closed set $\Omega' \neq \Omega$. Define $\chi: \Omega \rightarrow \{0, 1\}$ by $\chi(t) = 1$ iff $t \in \Omega'$, and put $(f, g, h) = f\bar{g}h(2\chi - 1)$, $f, g, h \in \mathcal{C}(\Omega)$. Then $(\mathcal{C}(\Omega), (\cdot, \cdot, \cdot), \|\cdot\|_{\sup})$ is a ternary C^* -ring, whose decomposition in the

sense of 3.1(1) is given by $\mathcal{C}(\Omega)_+ = \mathcal{C}(\Omega)\chi = \mathcal{C}(\Omega')$, $\mathcal{C}(\Omega)_- = \mathcal{C}(\Omega \setminus \Omega')$. Consequently, $Tf = f(2\chi - 1)$ for each $f \in \mathcal{C}(\Omega)$ and so $T \neq I$.

The problem of characterizing the ternary C^* -rings which yield $T = I$ remains open in the following. Let us mention two further points. Since, in view of Proposition 3.4, $\mathfrak{U}_b(\mathfrak{X})$ commutes with $\mathfrak{U}_a(\mathfrak{X})$ and hence, $T = 2P - 1$ belongs to the center of $M(\mathfrak{U})$ by Proposition 3.7, we see that $T \in \{I, -I\}$ if \mathfrak{U} is a factor. If \mathfrak{X} contains a "right unit," i.e., an element r such that $x = (x, r, r)$, $x \in \mathfrak{X}$, then $T = I$. Indeed, $x = x \cdot a(r, r)$ implies that $a(r, r)$ is a unit in \mathfrak{U} . Therefore, $r \in \mathfrak{X}_+$. Since $b(\mathfrak{X}_-, \mathfrak{X}_+) = \{0\}$ (cf. the proof of 3.7) we have that $x = (x, r, r) = b(x, r) \cdot r = 0$ for all $x \in \mathfrak{X}_-$.

4. THE REPRESENTATION THEOREM FOR TERNARY W^* -RINGS

A ternary W^* -ring $(\mathfrak{X}, (\cdot, \cdot, \cdot), \|\cdot\|)$ with a predual \mathfrak{X}_* and the weak* topology $\sigma = \sigma(\mathfrak{X}, \mathfrak{X}_*)$ will be called *normally isomorphic* (respectively, *normally anti-isomorphic*) to a TRO, if there are Hilbert spaces H and K and a linear isometry $U: \mathfrak{X} \rightarrow \mathcal{L}(H, K)$ which is a homeomorphism relative to σ and the weak-operator topology and which satisfies

$$U((x, y, z)) = U(x) U(y)^* U(z), \quad x, y, z \in \mathfrak{X}$$

(respectively, $U((x, y, z)) = -U(x) U(y)^* U(z)$, $x, y, z \in \mathfrak{X}$). Because of the weak* compactness of $D_1(\mathfrak{X})$ and the density theorem 1.4, $U(\mathfrak{X})$ is a weakly closed TRO.

The main result of this section consists now in the following:

4.1. REPRESENTATION THEOREM. *Let $(\mathfrak{X}, (\cdot, \cdot, \cdot), \|\cdot\|)$ be a ternary W^* -ring, $\mathfrak{X} = \mathfrak{X}_+ \oplus \mathfrak{X}_-$ its decomposition according to 3.1(1), and T the operator of 3.1(2). Then:*

(1) \mathfrak{X}_+ (respectively, \mathfrak{X}_-) is σ -closed and a ternary W^* -subring of \mathfrak{X} which is normally isomorphic (respectively, normally anti-isomorphic) to a TRO;

(2) T is σ -continuous, and the ternary W^* -ring $(\mathfrak{X}, T \circ (\cdot, \cdot, \cdot), \|\cdot\|)$ is normally isomorphic to a TRO.

In the proof we use the C^* -algebra \mathfrak{U} and the \mathfrak{U} -form α provided by Proposition 3.2. The essential step is Proposition 4.9 which states that the double centralizer algebra $M(\mathfrak{U})$ of \mathfrak{U} is a W^* -algebra and that \mathfrak{X} , endowed with the positive form $\alpha_T: (x, y) \mapsto \alpha(Tx, y)$, is a self-dual Hilbert $M(\mathfrak{U})$ -module. Theorem 4.1 is then a consequence of Theorem 2.8. Subsequently, we obtain that $M(\mathfrak{U})_*$ is equal to a quotient of $\mathfrak{X} \hat{\otimes}_\gamma \mathfrak{X}_*$ and that \mathfrak{X}_*

coincides with $\mathfrak{X}_c \otimes M(\mathfrak{U})_*$. In particular, \mathfrak{X}_* is uniquely determined up to isomorphism.

To establish Proposition 4.9 we proceed as follows: Replacing \mathfrak{a} by \mathfrak{a}_T we suppose that $(\mathfrak{X}, \mathfrak{a})$ is a Hilbert \mathfrak{U} -module. Let $(\mathfrak{X}'', \mathfrak{a})$ denote its self-dual completion and σ'' the weak topology of \mathfrak{X}'' (cf. Proposition 2.2 and Definition 2.3). Applying Theorem 2.6 we regard \mathfrak{X} as a norm closed TRO between H and K with weak closure \mathfrak{X}'' such that $[\mathfrak{X}]_H$ (respectively, $\{\mathfrak{X}\}_H$) is equal to \mathfrak{U} (respectively, \mathfrak{U}^{**}) and σ'' and $\sigma(\mathfrak{U}^{**}, \mathfrak{U}^*)$ are just the weak-operator topologies. Moreover, it is easy to see that $[\mathfrak{X}]_K$ coincides with the C^* -algebra \mathfrak{B} generated by $\mathfrak{b}(\mathfrak{X}, \mathfrak{X})$ (cf. Proposition 3.4). Our aim is now to construct two weak* continuous projections $\Phi: \mathfrak{X}'' \rightarrow \mathfrak{X}$ and $\Psi: \mathfrak{U}^{**} \rightarrow M(\mathfrak{U})$ of norm one which, for a central projection $e \in \mathfrak{U}^{**}$ satisfy

$$\text{kernel}(\Phi) = \mathfrak{X}''(1 - e) \quad \text{and} \quad \text{kernel}(\Psi) = \mathfrak{U}^{**}(1 - e).$$

Here, we use the fact that $M(\mathfrak{U})$ is a C^* -subalgebra of \mathfrak{U}^{**} . The idea of proof in the following is taken from [9, Lemma 1.7.6]:

4.2. LEMMA. *If $E \in [\mathfrak{X}]_H$ and $E' \in [\mathfrak{X}]_K$ are self-adjoint projections, then $\mathfrak{X} \cdot E$, $E' \cdot \mathfrak{X}$ are $(1 - E') \cdot \mathfrak{X}$ are σ -closed in \mathfrak{X} .*

Proof. Let $\{x_\alpha\}_\alpha \subset D_1(\mathfrak{X}) \cap \mathfrak{X} \cdot E$ be a net which tends to $x \in \mathfrak{X}$ in σ . We have to show that $x = x \cdot E$.

Suppose by way of contradiction that $x' = x \cdot (1 - E) \neq 0$. In view of Proposition 1.2, the partial isometry r of the polar decomposition of x' belongs to \mathfrak{X}'' . According to Proposition 1.4, there is a net $\{y_\beta\}_\beta \subset D_1(\mathfrak{X})$ which tends weakly to r . Since $\tau \mapsto \tau \cdot (1 - E)$ is weakly continuous on \mathfrak{X}'' , we may assume that $y_\beta = y_\beta \cdot (1 - E)$ for all β . By Notations 1.1 and Proposition 1.2, x' is positive in the W^* -algebra $\mathfrak{X}''_{(r)}$. Hence, there exists $\lambda > 0$ satisfying

$$\lambda + n \leq \|x' + nr\| = \liminf_{\beta} \|x' + ny_\beta\|, \quad n \in \mathbb{N}.$$

On the other hand, for each $y \in D_1(\mathfrak{X})$ with $y = y \cdot (1 - E)$ we have

$$\|x_\alpha + ny\| = \|\mathfrak{b}(x_\alpha, x_\alpha) + n^2 \mathfrak{b}(y, y)\|^{1/2} \leq (1 + n^2)^{1/2}, \quad n \in \mathbb{N}, \quad \text{all } \alpha,$$

and thus, together with $x + ny_\beta = \sigma - \lim_{\alpha} (x_\alpha + ny_\beta)$,

$$\|x' + ny_\beta\| = \|(x + ny_\beta) \cdot (1 - E)\| \leq (1 + n^2)^{1/2}, \quad n \in \mathbb{N}, \quad \text{all } \beta,$$

which leads to $\lambda + n \leq (1 + n^2)^{1/2}$, $n \in \mathbb{N}$, a contradiction.

The proof in the case of E' and $1 - E'$ is the same.

4.3. PROPOSITION. *Let $f \in \mathfrak{X}_*$. Then there exist a partial isometry $r \in \mathfrak{X}$ and positive functionals $\varphi \in \mathfrak{U}^*$, $\psi \in \mathfrak{B}^*$ such that*

- (1) $\varphi(a(x, r)) = f(x) = \psi(b(x, r)), x \in \mathfrak{X}$,
- (2) $f(r \cdot a) = \varphi(a), a \in \mathfrak{U}$,
- (3) $f(b \cdot r) = \psi(b), b \in \mathfrak{B}$.

Proof. We can suppose that $\|f\| = 1$. Since $D_1(\mathfrak{X})$ is compact in σ , there is an extreme point r of $D_1(\mathfrak{X})$ with $f(r) = 1$. In view of Lemma 1.3, r is a partial isometry. Now define $\varphi \in \mathfrak{U}^*$ by $\varphi(a) = f(r \cdot a)$. From $\varphi(a(r, r)) = f(r \cdot a(r, r)) = f(r) = 1$ we conclude that $\|\varphi\| = 1$. Hence, due to [9, Proposition 1.5.2], φ is positive. Moreover, $\varphi(a(x, r)) = f(r \cdot a(x, r)) = f(b(r, r) \cdot x)$, $x \in \mathfrak{X}$. To see that $f(x) = f(b(r, r) \cdot x)$, we argue by contradiction. Put $E' = b(r, r)$ and suppose that $f(x) \neq 0$ for some $x \in (I - E') \cdot \mathfrak{X}$. By Lemma 4.2, $(I - E') \cdot \mathfrak{X}$ is σ -closed and hence, a ternary W^* -ring. As above, there is a partial isometry $r' \in (I - E') \cdot \mathfrak{X}$ with $f(r') > 0$. Thus,

$$\begin{aligned} (1 + n^2)^{1/2} &\geq \|n^2 a(r, r) + a(r', r')\|^{1/2} = \|a(nr + r', nr + r')\|^{1/2} \\ &= \|nr + r'\| \geq |f(nr + r')| = n + f(r'), \quad n \in \mathbb{N}, \end{aligned}$$

a contradiction. The functional ψ is found similarly.

4.4. LEMMA. *Each $f \in \mathfrak{X}_*$ has a unique extension to a weakly continuous functional f'' on \mathfrak{X}'' . The adjoint $\Phi: \mathfrak{X}'' \rightarrow \mathfrak{X}$ of $f \mapsto f''$ is a linear, σ'' - σ -continuous projection of norm one.*

Proof. Let $f \in \mathfrak{X}_*$ and write $f = \varphi(a(\cdot, r))$ in accordance with 4.3(1), (2). Define f'' on \mathfrak{X}'' by $f''(\tau) = \varphi(a(\tau, r))$. Then f'' is a weakly continuous functional on \mathfrak{X}'' , because $a(\tau, r) = \tau(r)^*$ for all $\tau \in \mathfrak{X}''$ by 2.2(4). Obviously, f'' is an extension of f . Since, because of Proposition 1.4, $D_1(\mathfrak{X})$ is weakly dense in $D_1(\mathfrak{X}'')$ the extension is unique and satisfies $\|f\| = \|f''\|$.

We are now going to derive the module properties of Φ and to show that the kernel of Φ is a two-sided ideal in the sense of Lemma 1.5. First, we look at the behaviour of Φ on C^* -algebras which are associated with partial isometries in \mathfrak{X} .

4.5. LEMMA. *Let $r \in \mathfrak{X}$ be a partial isometry and $\mathfrak{X}_{(r)}$ (respectively, $\mathfrak{X}_{(r)}''$) the C^* -algebra (respectively, W^* -algebra) associated with r in \mathfrak{X} (respectively, \mathfrak{X}'') according to Notations 1.1. Then, $\Phi(\mathfrak{X}_{(r)}'') = \mathfrak{X}_{(r)}$. In particular, $\mathfrak{X}_{(r)}$ is a W^* -algebra with predual $\{f|_{\mathfrak{X}_{(r)}} | f \in \mathfrak{X}_*\}$.*

Proof. We start with the following claim:

Claim 4.5.1. Let $x \in \mathfrak{X}$. If $f(x) \geq 0$ for all $f \in \mathfrak{X}^*$ whose restriction to $\mathfrak{X}_{(r)}$ is a positive functional, then x is positive in $\mathfrak{X}_{(r)}$.

By the Hahn–Banach theorem, the state space of $\mathfrak{X}_{(r)}$ is equal to $\{f|_{\mathfrak{X}_{(r)}} | f \in \mathfrak{X}^*, f(r) = 1 = \|f\|\}$. Hence, it suffices to show that $x \in \mathfrak{X}_{(r)}$. Suppose that $x' = (1 - E') \cdot x \neq 0$, where E' denotes the range projection of r . Then there is $f \in \mathfrak{X}^*$ with $f(x') = -1$. Since the functional $g: y \mapsto f(y - E' \cdot y)$ vanishes on $\mathfrak{X}_{(r)}$ we obtain $g(x) \geq 0$ which contradicts $g(x) = f(x') = -1$. Similarly, $x = x \cdot a(r, r)$.

It follows that $\Phi(\mathfrak{X}_{(r)}'') \subset \mathfrak{X}_{(r)}$. Indeed, let $\tau \in \mathfrak{X}_{(r)}''$ be positive and fix $f \in \mathfrak{X}^*$ satisfying $f(r) = 1 = \|f\|$. If g denotes $f \circ \Phi$ restricted to $\mathfrak{X}_{(r)}$, then $g(r) = f(r) = 1$ and $\|g\| \leq 1$. Hence, g is a state. Therefore, $g(\tau) \geq 0$ and so, $f(\Phi\tau) \geq 0$. By Claim 4.5.1, $\Phi(\tau) \in \mathfrak{X}_{(r)}$.

Observe next that $\mathfrak{X}_{(r)}''$ is weakly closed in \mathfrak{X}'' . Since $D_1(\mathfrak{X}_{(r)}) = \Phi(D_1(\mathfrak{X}_{(r)}''))$ this ball is compact in σ and thus, $\mathfrak{X}_{(r)}$ is σ -closed.

4.6. LEMMA. For all $A \in M(\mathfrak{A})$ and $b \in \mathfrak{B}$, the mappings $x \mapsto x \cdot A$ and $x \mapsto b \cdot x$ are σ -continuous on \mathfrak{X} .

Proof. It suffices to show that $x \mapsto x \cdot U$ is σ -continuous for each unitary $U \in M(\mathfrak{A})$. So, let $\{x_\alpha\}_\alpha \subset \mathfrak{X}$ be a net tending in σ to zero and fix $f \in \mathfrak{X}_*$. We must prove that $\lim_\alpha f(x_\alpha \cdot U) = 0$.

Write $f = \psi(b(\cdot, r))$ according to Proposition 4.3. Put $E' = b(r, r)$ and $r' = r \cdot U^*$. Then r' is a partial isometry with range projection E' . Define $\mathfrak{C} = E'\mathfrak{B}E'$. In view of Proposition 1.2, a $*$ -isomorphism of \mathfrak{C} onto $\mathfrak{X}_{(r)}$ (respectively, $\mathfrak{X}_{(r')}$) is given by $b \mapsto b \cdot r$ (respectively, $b \mapsto b \cdot r'$). Using Lemma 4.5 we obtain that \mathfrak{C} is a W^* -algebra with predual $\{b \mapsto g(b \cdot r) | g \in \mathfrak{X}_*\} = \{b \mapsto g(b \cdot r') | g \in \mathfrak{X}_*\}$. Hence, ψ is normal on \mathfrak{C} and so, $\psi(b) = h(b \cdot r')$, $b \in \mathfrak{C}$ for some $h \in \mathfrak{X}_*$. By Lemma 4.2, the mappings $x \mapsto E' \cdot x$ and $x \mapsto x \cdot a(r', r')$ are σ -continuous on \mathfrak{X} . Thus,

$$\begin{aligned} \lim_\alpha f(x_\alpha \cdot U) &= \lim_\alpha \psi(b(x_\alpha \cdot U, r)) = \lim_\alpha \psi(b(E' \cdot x_\alpha, r')) \\ &= \lim_\alpha h(b(E' \cdot x_\alpha, r') \cdot r') = \lim_\alpha h(E' \cdot x_\alpha \cdot a(r', r')) = 0. \end{aligned}$$

4.7. COROLLARY. Let Φ be the projection defined in Lemma 4.4. Then

(1) $\Phi(\tau \cdot A) = \Phi(\tau) \cdot A$ and $\Phi(b \cdot \tau) = b \cdot \Phi(\tau)$ for all $\tau \in \mathfrak{X}''$, $A \in M(\mathfrak{A})$ and $b \in \mathfrak{B}$.

(2) The kernel of Φ , $\ker(\Phi)$, is of the form $\ker(\Phi) = \mathfrak{X}'' \cdot (I - e)$, where e is a (uniquely determined) central projection in \mathfrak{A}^{**} .

Proof. Let $\tau \in \mathfrak{X}''$. Then τ is the weak limit of a net $\{x_\alpha\}_\alpha \subset \mathfrak{X}$, and

$\{x_\alpha \cdot A\}_\alpha$ tends weakly to $\tau \cdot A$ for each $A \in M(\mathfrak{A})$. Using $\Phi(\tau) = \sigma - \lim_\alpha x_\alpha$ and Lemma 4.6 we obtain

$$\Phi(\tau \cdot A) = \sigma - \lim_\alpha \Phi(x_\alpha \cdot A) = \sigma - \lim_\alpha x_\alpha \cdot A = \Phi(\tau) \cdot A, \quad A \in M(\mathfrak{A}).$$

The second statement in (1) follows similarly. Now, (1) implies that $\ker(\Phi) \cdot \mathfrak{A} \cup \mathfrak{B} \cdot \ker(\Phi) \subset \ker(\Phi)$. By 1.5, $\ker(\Phi) = \mathfrak{X}'' \cdot (I - e)$ with a unique central projection $e \in \{\mathfrak{X}\}_H = \mathfrak{A}^{**}$.

4.8. LEMMA. *$M(\mathfrak{A})$ is a W^* -algebra. In particular, there exists a linear weak* continuous projection $\Psi: \mathfrak{A}^{**} \rightarrow M(\mathfrak{A})$ such that*

- (1) $\Psi(a(\tau, x)) = a(\Phi(\tau), x)$, $\tau \in \mathfrak{X}'', x \in \mathfrak{X}$;
- (2) $\ker(\Psi) = \mathfrak{A}^{**}(I - e)$, where e is the projection in 4.7(2).

Proof. Let $A \in \mathfrak{A}^{**}$. By 4.7(1), $\Phi_A: x \mapsto \Phi(x \cdot A)$ is a \mathfrak{B} -module homomorphism on \mathfrak{X} . We are going to show that $\Phi_A \in \mathfrak{U}_b(\mathfrak{X})$ and then define $\Psi(A) = \Phi_A$.

To this end we write Ω for the set of all bounded \mathfrak{B} -module homomorphisms on \mathfrak{X} . Observe that $M(\mathfrak{A}) \subset \Omega$ by 3.4(4). If Γ denotes the weak topology on Ω generated by all functionals $S \mapsto f(Sx)$, $f \in \mathfrak{X}_*$, $x \in \mathfrak{X}$, then we have:

LEMMA 4.8.1. *a is separately $(\sigma \times \sigma)$ - Γ -continuous.*

Proof. If $\{x_\alpha\}_\alpha \subset \mathfrak{X}$ is a net tending to $x \in \mathfrak{X}$ in σ , then Lemma 4.6 shows

$$\begin{aligned} \lim_\alpha f(y \cdot a(x_\alpha, z)) &= \lim_\alpha f(b(y, z) \cdot x_\alpha) = f(b(y, z) \cdot x) \\ &= f(y \cdot a(x, z)), \quad f \in \mathfrak{X}_*, \quad y, z \in \mathfrak{X}, \end{aligned}$$

and thus, $a(x, z) = \Gamma\text{-}\lim_\alpha a(x_\alpha, z)$. On the other hand, with $f = \varphi(a(\cdot, r))$ according to 4.3(1), (2) we conclude that

$$\begin{aligned} f(y \cdot a(z, x_\alpha)) &= \varphi(a(y, r) a(z, x_\alpha)) = \overline{\varphi(a(x_\alpha, z) a(r, y))} \\ &= \overline{\varphi(a(r, r) a(x_\alpha, z) a(r, y))} = \overline{\varphi(a(r \cdot a(x_\alpha, z) a(r, y), r))} \\ &= \overline{\varphi(a(b(r, z) \cdot x_\alpha \cdot a(r, y), r))} = \overline{f(b(r, z) \cdot x_\alpha \cdot a(r, y))} \\ &\xrightarrow{\alpha} \overline{f(b(r, z) \cdot x \cdot a(r, y))} = \dots = f(y \cdot a(z, x)). \end{aligned}$$

Hence, $a(z, x) = \Gamma\text{-}\lim_\alpha a(z, x_\alpha)$, $z \in \mathfrak{X}$, proving Lemma 4.8.1.

Now, let A be the weak*-limit of a net $\{a_\alpha\}_\alpha \subset \mathfrak{A}$. Then

$$a(z, \Phi_A(y)) = \Gamma\text{-}\lim_\alpha a(z, y \cdot a_\alpha^*), \quad y, z \in \mathfrak{X},$$

which follows from Lemma 4.8.1, because $y \cdot A^* = \sigma''\text{-}\lim_{\alpha} y \cdot a_{\alpha}^*$ and so, $\Phi_{A^*}(y) = \sigma\text{-}\lim_{\alpha} y \cdot a_{\alpha}^*$. Therefore, we obtain

$$\begin{aligned} b(\Phi_A(x), y) \cdot z &= \Phi(x \cdot A) \cdot a(z, y) = \Phi(x \cdot A a(z, y)) \\ &= \sigma\text{-}\lim_{\alpha} x \cdot a_{\alpha} a(z, y) = \sigma\text{-}\lim_{\alpha} x \cdot a(z, y \cdot a_{\alpha}^*) \\ &= x \cdot a(z, \Phi_{A^*}(y)) = b(x, \Phi_{A^*}(y)) \cdot z, \end{aligned}$$

and consequently, $\Phi_A \in \mathfrak{U}_b(\mathfrak{X})$ and $(\Phi_A)^* = \Phi_{A^*}$.

Obviously, $\Psi: A \rightarrow \Phi_A$ is a $*$ -anti-homomorphism of \mathfrak{U}^{**} onto $\mathfrak{U}_b(\mathfrak{X}) = M(\mathfrak{U})_{\sim}$. Because of 4.7(1), Ψ is a projection. Moreover,

$$\ker(\Psi) = \{A \in \mathfrak{U}^{**} \mid \mathfrak{X} \cdot A \subset \ker(\Phi)\} = \mathfrak{U}^{**}(I - e).$$

Therefore, $M(\mathfrak{U})$ is a W^* -algebra which is isomorphic to $\mathfrak{U}^{**}e$, and Ψ is weak $*$ continuous. Finally, suppose that $\tau = \sigma''\text{-}\lim_{\alpha} x_{\alpha}$ for a net $\{x_{\alpha}\}_{\alpha} \subset \mathfrak{X}$. Then (1) follows from

$$\begin{aligned} y \cdot \Psi(a(\tau, x)) &= \sigma\text{-}\lim_{\alpha} \Phi(y \cdot a(x_{\alpha}, x)) = \sigma\text{-}\lim_{\alpha} b(y, x) \cdot x_{\alpha} \\ &= b(y, x) \cdot \Phi(\tau) = y \cdot a(\Phi(\tau), x). \end{aligned}$$

4.9. PROPOSITION. *Let $(\mathfrak{X}(\cdot, \cdot, \cdot), \|\cdot\|)$ be a ternary W^* -ring and $T: \mathfrak{X} \rightarrow \mathfrak{X}$ the operator belonging to it by 3.1(2). Then $M(\mathfrak{U})$ is a W^* -algebra, and (\mathfrak{X}, a_T) is a self-dual Hilbert $M(\mathfrak{U})$ -module, a_T being defined by $a_T(x, y) = a(Tx, y)$.*

Proof. Let $\tau: \mathfrak{X} \rightarrow M(\mathfrak{U})$ be a bounded $M(\mathfrak{U})$ -module homomorphism. Then $\tau_1 = \tau \circ \Phi$ is in \mathfrak{X}'' , and 4.8(1) shows that

$$\tau(x) = \tau_1(x) = \Psi(\tau_1(x)^*)^* = \Psi(a_T(\tau_1, x))^* = a_T(x, \Phi(\tau_1)), \quad x \in \mathfrak{X}.$$

Hence, (\mathfrak{X}, a_T) is self-dual over $M(\mathfrak{U})$.

Proof of 4.1. By virtue of Proposition 3.7, the projection $P: \mathfrak{X} \rightarrow \mathfrak{X}_+$ belongs to the center of $M(\mathfrak{U})$. By Lemma 4.6, P is σ -continuous. Therefore, \mathfrak{X}_+ and \mathfrak{X}_- are σ -closed, and $T = 2P - I$ is σ -continuous. Because of Proposition 4.9, $(\mathfrak{X}_{\pm}, a \mid \mathfrak{X}_{\pm} \times \mathfrak{X}_{\pm})$ and (\mathfrak{X}, a_T) are self-dual Hilbert modules over the W^* -algebra $M(\mathfrak{U})$. From Theorem 2.8 we obtain the desired isomorphisms.

4.10. COROLLARY. *Let $(\mathfrak{X}, (\cdot, \cdot, \cdot), \|\cdot\|)$ be a ternary W^* -ring and $\mathfrak{U}, \mathfrak{B}$ the C^* -algebras as above. Then $M(\mathfrak{U})_{\sim}$ is the set of all bounded $M(\mathfrak{B})$ -module homomorphisms on \mathfrak{X} . In particular, $M(\mathfrak{U})$ is the dual space of the quotient $\mathfrak{X} \otimes_{M(\mathfrak{B})} \mathfrak{X}_*$ of $\mathfrak{X} \hat{\otimes}_{\gamma} \mathfrak{X}_*$.*

Proof. Define $b_T: \mathfrak{X} \times \mathfrak{X} \rightarrow M(\mathfrak{B})$ by $b_T(x, y) = b(Tx, y)$. If we apply the same line of reasoning by which Proposition 4.9 has been established to $(\mathfrak{X}, \mathfrak{B}, b_T)$ instead of $(\mathfrak{X}, \mathfrak{A}, a_T)$ we obtain that (\mathfrak{X}, b_T) is a self-dual (left) Hilbert module over the W^* -algebra $M(\mathfrak{B})$. In view of [5, Corollary 3.5], the set of all bounded $M(\mathfrak{B})$ -module homomorphisms is equal to $\mathfrak{A}_{b_T}(\mathfrak{X})$. Now, $\mathfrak{A}_{b_T}(\mathfrak{X}) = \mathfrak{A}_b(\mathfrak{X})$ and, by 3.4(3), $\mathfrak{A}_b(\mathfrak{X}) = M(\mathfrak{A}_\sim) = M(\mathfrak{A})_\sim$, and the first assertion follows. For the proof of the second statement, we observe that \mathfrak{X}_* is a left Banach $M(\mathfrak{B})$ -module. Hence, [7, Corollary 2.13] shows that $(\mathfrak{X} \otimes_{M(\mathfrak{B})} \mathfrak{X}_*)^* = M(\mathfrak{A})$.

4.11. COROLLARY. *Let $(\mathfrak{X}, (\cdot, \cdot, \cdot), \|\cdot\|)$ be a ternary W^* -ring. Then the predual of \mathfrak{X} consists precisely of all functionals $x \mapsto \varphi(a_T(x, y))$, $y \in \mathfrak{X}$, $\varphi \in M(\mathfrak{A})_*$ and is therefore uniquely determined up to isomorphism.*

Proof. Let f be a weak* continuous functional on \mathfrak{X} . By Proposition 4.3, there exist $r \in \mathfrak{X}$ and $\varphi \in M(\mathfrak{A})^*$ such that $f = \varphi(a_T(\cdot, r))$ and $\varphi = f(r \cdot)$. In particular, φ is Γ -continuous on $M(\mathfrak{A})$, where Γ denotes the topology defined in the proof of Lemma 4.8. Since the projection $\Psi: \mathfrak{A}^{**} \rightarrow M(\mathfrak{A})$ is weak*- Γ -continuous we conclude that $\varphi \circ \Psi \in \mathfrak{A}^*$. Consequently, $\varphi \in M(\mathfrak{A})_*$ so that f has the desired representation.

On the other hand, fix $y \in \mathfrak{X}$, $\varphi \in M(\mathfrak{A})_*$ and put $g = \varphi(a_T(\cdot, y))$. Then Corollary 4.10 shows the existence of sequences $\{x_n\}_n \subset \mathfrak{X}$, $\{f_n\}_n \subset \mathfrak{X}_*$ satisfying $\sum_n \|x_n\| \|f_n\| < \infty$ and $\varphi(A) = \sum_n f_n(x_n \cdot A)$, $A \in M(\mathfrak{A})$. Hence, the functional $f: x \mapsto \sum_n f_n(b_T(x_n, y) \cdot x)$ belongs to \mathfrak{X}_* . Therefore, $g \in \mathfrak{X}_*$, because $g(x) = \sum_n f_n(x_n \cdot a_T(x, y)) = f(x)$.

In our former considerations we have already used the fact that each self-dual Hilbert module $(\mathfrak{H}, \langle \cdot | \cdot \rangle)$ over a W^* -algebra \mathfrak{A} is a dual space. Hence, $(\mathfrak{H}, (\cdot, \cdot, \cdot), \|\cdot\|_{\mathfrak{H}})$ is a ternary W^* -ring, (\cdot, \cdot, \cdot) given by $(x, y, z) = x \cdot \langle z | y \rangle$. Now, Corollary 4.11 implies that the canonical predual $\mathfrak{H}_c \otimes_{\mathfrak{A}} \mathfrak{A}_*$ coincides with $\mathfrak{H}_c \otimes \mathfrak{A}_*$ and is uniquely determined up to isomorphism. Our final example includes the converse of Theorem 2.8.

4.12. EXAMPLE. Let $\mathfrak{R} \subset \mathcal{L}(H, K)$ be a weakly closed TRO. Then $(\mathfrak{R}, \langle \cdot | \cdot \rangle)$ is a self-dual Hilbert $\{\mathfrak{R}\}_H$ -module, where $\langle A | B \rangle = B^*A$. In particular, $(\mathfrak{R}, (\cdot, \cdot, \cdot), \|\cdot\|)$ is a ternary W^* -ring with (\cdot, \cdot, \cdot) given by $(A, B, C) = AB^*C$ and with unique predual $\mathfrak{R}_c \otimes (\{\mathfrak{R}\}_H)_*$.

Proof. To show self-duality, let $\tau: \mathfrak{R} \rightarrow \{\mathfrak{R}\}_H$ be a bounded $\{\mathfrak{R}\}_H$ -module homomorphism. Since $(A, f) \mapsto f(\tau(A)^*)$ is continuous on $\mathfrak{R}_c \times (\{\mathfrak{R}\}_H)_*$ there exists an η in the dual space of the projective tensor product $\mathfrak{R}_c \hat{\otimes}_{\gamma} (\{\mathfrak{R}\}_H)_*$ satisfying $\eta(A \otimes f) = f(\tau(A)^*)$, $A \in \mathfrak{R}$. Since

$$\eta((AX) \otimes f - A \otimes (f \cdot X)) = f(\tau(AX)^*) - f(X^* \tau(A)^*) = 0$$

we may assume that $\eta \in \mathfrak{R}$ which in turn yields

$$f(\tau(A)^*) = \eta(A \otimes f) = f(A^*\eta), \quad A \in \mathfrak{R}, \quad f \in (\{\mathfrak{R}\}_H)_*$$

and therefore, $\tau = \langle \cdot | \eta \rangle$.

ACKNOWLEDGMENTS

I should like to acknowledge my sincere thanks to Professor D. Kölzow for his help and encouragement during the preparation of this work and to Professor E. Albrecht for some valuable comments on the subject.

REFERENCES

1. R. C. BUSBY, Double centralizers of C^* -algebras, *Trans. Amer. Math. Soc.* **132** (1968), 79–99.
2. M. R. HESTENES, A ternary algebra with applications to matrices and linear transformations, *Arch. Rational Mech. Anal.* **11** (1962), 138–194.
3. M. R. HESTENES, Relative self-adjoint operators in Hilbert space, *Pacific J. Math.* **11** (1961), 1315–1357.
4. I. KAPLANSKY, Modules over operator algebras, *Amer. J. Math.* **75** (1953), 839–858.
5. W. L. PASCHKE, Inner product modules over B^* -algebras, *Trans. Amer. Math. Soc.* **182** (1973), 443–468.
6. W. L. PASCHKE, Inner product modules arising from compact automorphism groups of von Neumann algebras, *Trans. Amer. Math. Soc.* **224** (1976), 87–102.
7. M. A. RIEFFEL, Induced Banach representations of Banach algebras and locally compact groups, *J. Funct. Anal.* **1** (1967), 443–491.
8. M. A. RIEFFEL, Induced representations of C^* -algebras, *Advan. in Math.* **13** (1974), 176–257.
9. S. SAKAI, C^* -algebras and W^* -algebras, Springer, Berlin/New York, 1971.
10. M. F. SMILEY, An introduction to Hestenes ternary rings, *Amer. Math. Monthly* **76** (1969), 245–248.
11. R. A. STEPHENSON, Jacobson structure theory for Hestenes ternary rings, *Trans. Amer. Math. Soc.* **177** (1973), 91–98.
12. D. C. TAYLOR, The strict topology for double centralizer algebras, *Trans. Amer. Math. Soc.* **150** (1970), 633–643.
13. J. H. WILLIAMSON (Ed.), "Algebras in Analysis," Academic Press, New York, 1975.